

RC
268.5
U55
no.35
1978

National Cancer Institute
CARCINOGENESIS
Technical Report Series
No. 35
1978

**BIOASSAY OF
METHOXYCHLOR
FOR POSSIBLE CARCINOGENICITY**

CAS No. 72-43-5

NCI-CG-TR-35

U.S. DEPARTMENT OF HEALTH, EDUCATION, AND WELFARE
Public Health Service
National Institutes of Health



Library
National Institutes of Health
Bethesda, Maryland 20814

U. S. National Cancer Institute

BIOASSAY OF
METHOXYCHLOR
FOR POSSIBLE CARCINOGENICITY

Carcinogenesis Testing Program
Division of Cancer Cause and Prevention
National Cancer Institute
National Institutes of Health
Bethesda, Maryland 20014

U.S. DEPARTMENT OF HEALTH, EDUCATION, AND WELFARE
Public Health Service
National Institutes of Health

DHEW Publication No. (NIH) 78-835

PC

268.5

2155

no. 35

1978

REPORT ON THE BIOASSAY OF METHOXYCHLOR
FOR POSSIBLE CARCINOGENICITY

CARCINOGENESIS TESTING PROGRAM
DIVISION OF CANCER CAUSE AND PREVENTION
NATIONAL CANCER INSTITUTE, NATIONAL INSTITUTES OF HEALTH

CONTRIBUTORS: This report presents the results of the bioassay of methoxychlor conducted for the Carcinogenesis Testing Program, Division of Cancer Cause and Prevention, National Cancer Institute (NCI), National Institutes of Health, Bethesda, Maryland. This bioassay was conducted by Hazleton Laboratories America, Inc., Vienna, Virginia, initially under direct contract to the NCI and currently under a sub-contract to Tracor Jitco, Inc., prime contractor for the NCI Carcinogenesis Bioassay Program.

The experimental design was determined by the NCI Project Officers, Dr. J. H. Weisburger (1,2) and Dr. E. K. Weisburger (1). The principal investigators for the contract were Dr. M. B. Powers (3), Dr. R. W. Voelker (3), Dr. W. A. Olson (3,4) and Dr. W. M. Weatherholtz (3). Chemical analysis was performed by Dr. C. L. Guyton (3,5); the technical supervisor of animal treatment and observation was Ms. K. J. Petrovics (3).

Histopathologic examinations were performed by Dr. R. H. Habermann (3) and reviewed by Dr. R. W. Voelker (3) at the Hazleton Laboratories America, Inc., and the diagnoses included in this report represent the interpretation of these pathologists. Histopathology findings and reports were reviewed by Dr. R. L. Schueler (6).

Compilation of individual animal survival, pathology, and summary tables was performed by EG&G Mason Research Institute (7); the statistical analysis was performed by Dr. J. R. Joiner (6), and Mr. W. W. Belew (8), using methods selected for the Bioassay Program by Dr. J. J. Gart (9).

This report was prepared at METREK, a Division of The MITRE Corporation (8) under the direction of the NCI. Those responsible for this report at METREK are the project coordinator, Dr. L. W. Thomas (8), the task leader, Dr. M. R. Kornreich (8), and the senior biologist, Ms. P. Walker (8). The final report was reviewed by members of the participating organizations.

The statistical analysis was reviewed by one or more members of the Mathematical Statistics and Applied Mathematics Section of the NCI: Dr. J. J. Gart (9), Mr. J. Nam (9), Dr. H. M. Pettigrew (9), and Dr. R. E. Tarone (9).

The following other scientists at the National Cancer Institute were responsible for evaluating the bioassay experiment, interpreting the results, and reporting the findings: Dr. K. C. Chu (1), Dr. C. Cueto, Jr. (1), Dr. J. F. Douglas (1), Dr. D. G. Goodman (1), Dr. R. A. Griesemer (1), Dr. H. Milman (1), Dr. T. W. Orme (1), Dr. R. A. Squire (1,10), and Dr. J. M. Ward (1).

-
1. Carcinogenesis Testing Program, Division of Cancer Cause and Prevention, National Cancer Institute, National Institutes of Health, Bethesda, Maryland.
 2. Now with the Naylor Dana Institute for Disease Prevention, American Health Foundation, Hammon House Road, Valhalla, New York.
 3. Hazleton Laboratories America, Inc., 9200 Leesburg Turnpike, Vienna, Virginia.
 4. Now with the Center for Regulatory Services, 2347 Paddock Lane, Reston, Virginia.
 5. Now with Rhodia, Inc., 23 Belmont Drive, Somerset, New Jersey.
 6. Tracor Jitco, Inc., 1776 East Jefferson Street, Rockville, Maryland.
 7. EG&G Mason Research Institute, 1530 East Jefferson Street, Rockville, Maryland.
 8. The MITRE Corporation, METREK Division, 1820 Dolley Madison Boulevard, McLean, Virginia.
 9. Mathematical Statistics and Applied Mathematics Section, Biometry Branch, Field Studies and Statistics Program, Division of Cancer Cause and Prevention, National Cancer Institute, National Institutes of Health, Bethesda, Maryland.
 10. Now with the Division of Comparative Medicine, Johns Hopkins University, School of Medicine, Traylor Building, Baltimore, Maryland.

SUMMARY

A bioassay for possible carcinogenicity of technical-grade methoxychlor was conducted using Osborne-Mendel rats and B6C3F1 mice. Methoxychlor was administered in the feed, at either of two concentrations, to groups of 50 male and 50 female animals of each species. For each species, 20 animals of each sex were placed on test as controls. The time-weighted average high and low dietary concentrations of methoxychlor were, respectively, 845 and 448 ppm for male rats, 1385 and 750 ppm for female rats, 3491 and 1746 ppm for male mice, and 1994 and 997 ppm for female mice. After a treatment period of 78 weeks, the rat groups were observed for an additional 34 weeks and the mouse groups for an additional 15 weeks. A dose-related mean group body weight depression was observed in both rats and mice, but no effect on survival was detected.

Under the conditions of this study, methoxychlor was not found to be carcinogenic in Osborne-Mendel rats or B6C3F1 mice of either sex.

TABLE OF CONTENTS

	<u>Page</u>
I. INTRODUCTION	1
II. MATERIALS AND METHODS	5
A. Chemicals	5
B. Dietary Preparation	5
C. Animals	5
D. Animal Maintenance	6
E. Selection of Initial Concentrations	7
F. Experimental Design	8
G. Clinical and Histopathologic Examinations	11
H. Data Recording and Statistical Analyses	13
III. CHRONIC TESTING RESULTS: RATS	19
A. Body Weights and Clinical Observations	19
B. Survival	21
C. Pathology	21
D. Statistical Analyses of Results	23
IV. CHRONIC TESTING RESULTS: MICE	35
A. Body Weights and Clinical Observations	35
B. Survival	35
C. Pathology	38
D. Statistical Analyses of Results	38
V. DISCUSSION	43
VI. BIBLIOGRAPHY	45
APPENDIX A SUMMARY OF THE INCIDENCE OF NEOPLASMS IN RATS TREATED WITH METHOXYCHLOR	A-1
APPENDIX B SUMMARY OF THE INCIDENCE OF NEOPLASMS IN MICE TREATED WITH METHOXYCHLOR	B-1
APPENDIX C SUMMARY OF THE INCIDENCE OF NONNEOPLASTIC LESIONS IN RATS TREATED WITH METHOXYCHLOR	C-1
APPENDIX D SUMMARY OF THE INCIDENCE OF NONNEOPLASTIC LESIONS IN MICE TREATED WITH METHOXYCHLOR	D-1

LIST OF ILLUSTRATIONS

<u>Figure Number</u>		<u>Page</u>
1	GROWTH CURVES FOR METHOXYCHLOR CHRONIC STUDY RATS	20
2	SURVIVAL COMPARISONS OF METHOXYCHLOR CHRONIC STUDY RATS	22
3	GROWTH CURVES FOR METHOXYCHLOR CHRONIC STUDY MICE	36
4	SURVIVAL COMPARISONS OF METHOXYCHLOR CHRONIC STUDY MICE	37

LIST OF TABLES

<u>Table Number</u>		<u>Page</u>
1	DESIGN SUMMARY FOR OSBORNE-MENDEL RATS--METHOXYCHLOR FEEDING EXPERIMENT	9
2	DESIGN SUMMARY FOR B6C3F1 MICE--METHOXYCHLOR FEEDING EXPERIMENT	10
3	ANALYSES OF THE INCIDENCE OF PRIMARY TUMORS AT SPECIFIC SITES IN MALE RATS TREATED WITH METHOXYCHLOR	24
4	ANALYSES OF THE INCIDENCE OF PRIMARY TUMORS AT SPECIFIC SITES IN FEMALE RATS TREATED WITH METHOXYCHLOR	30
5	ANALYSES OF THE INCIDENCE OF PRIMARY TUMORS AT SPECIFIC SITES IN MALE MICE TREATED WITH METHOXYCHLOR	39
6	ANALYSES OF THE INCIDENCE OF PRIMARY TUMORS AT SPECIFIC SITES IN FEMALE MICE TREATED WITH METHOXYCHLOR	40
A1	SUMMARY OF THE INCIDENCE OF NEOPLASMS IN MALE RATS TREATED WITH METHOXYCHLOR	A-3

LIST OF TABLES (Concluded)

<u>Table Number</u>		<u>Page</u>
A2	SUMMARY OF THE INCIDENCE OF NEOPLASMS IN FEMALE RATS TREATED WITH METHOXYCHLOR	A-7
B1	SUMMARY OF THE INCIDENCE OF NEOPLASMS IN MALE MICE TREATED WITH METHOXYCHLOR	B-3
B2	SUMMARY OF THE INCIDENCE OF NEOPLASMS IN FEMALE MICE TREATED WITH METHOXYCHLOR	B-6
C1	SUMMARY OF THE INCIDENCE OF NONNEOPLASTIC LESIONS IN MALE RATS TREATED WITH METHOXYCHLOR	C-3
C2	SUMMARY OF THE INCIDENCE OF NONNEOPLASTIC LESIONS IN FEMALE RATS TREATED WITH METHOXYCHLOR	C-7
D1	SUMMARY OF THE INCIDENCE OF NONNEOPLASTIC LESIONS IN MALE MICE TREATED WITH METHOXYCHLOR	D-3
D2	SUMMARY OF THE INCIDENCE OF NONNEOPLASTIC LESIONS IN FEMALE MICE TREATED WITH METHOXYCHLOR	D-6



I. INTRODUCTION

Methoxychlor (NCI No. C00497), a synthetic organochlorine insecticide and structural analog of DDT, was one of several widely used agricultural pesticides selected for bioassay by the National Cancer Institute because of a lack of adequate chronic toxicity data. The suspect state of all DDT-related chemicals was an important additional factor in its selection.

The Chemical Abstracts Service (CAS) Ninth Collective Index

(1977) name for this compound is 1,1'-(2,2,2-trichloroethylidene)bis(4-methoxy)-benzene.* It is also known as 1,1,1-trichloro-2,2'-bis(p-methoxyphenyl)-ethane; dianisyl trichloroethane; dimethoxy DDT; and DMDT.

Methoxychlor is effective against a wide range of insects that attack fruits, vegetables, shade trees, home gardens, forage crops, and livestock. It is also used for the spray treatment of barns, grain storage bins, mushroom houses, dairies, and other agricultural buildings (Spencer, 1973; Brooks, 1974; Farm Chemicals Handbook, 1976). In 1971, the most recent year for which data are available, more than 3 million pounds of methoxychlor were used in the United States. Of this amount, approximately 2 million pounds were used to protect livestock and livestock buildings, with application to beef and dairy cattle accounting for 95 percent of livestock-related usage. The remaining 1 million pounds of methoxychlor were used to treat 725

*The CAS registry number is 72-43-5.

thousand acres of varied cropland. The major crop treated was alfalfa which accounted for 49 percent of crop-related methoxychlor usage and 66 percent of total treated acreage (Andrilenas, 1974). Although its activity range differs somewhat from that of DDT, methoxychlor is considered to be an excellent replacement for DDT for certain applications (Spencer, 1973; Farm Chemicals Handbook, 1976). It may also substitute in some cases for aldrin, dieldrin, chlordane and heptachlor, all of which have had their registrations cancelled (Ouellette and King, 1976).

According to the U.S. International Trade Commission (1977) 5.504×10^6 pounds of methoxychlor were produced in the United States in 1975, an almost 70 percent increase over 1974 production figures (3.248×10^6 pounds) (Fowler and Mahan, 1976).

The risk of exposure to methoxychlor is greatest for agricultural or farm workers engaged in pesticide application, although workers at pesticide formulating plants may also experience significant contact with the chemical. Since methoxychlor is relatively ineffective against soil organisms (Brooks, 1974), it is generally applied directly to crops via ground or aerial spraying. As a result, contamination of the atmosphere over treated lands and surrounding areas, with possible inhalational exposure of persons residing in those areas, may occur.

Methoxychlor possesses long residual insecticidal activity against many species (Farm Chemicals Handbook, 1976) probably as a

result of its low volatility and relative resistance to oxidation, heat, moisture and ultraviolet irradiation (Brooks, 1974). Consequently, residues may persist on treated crops for a length of time sufficient to permit low-level exposure among the general population. As an example, 5.3 ppm of methoxychlor were detected on alfalfa 33 days following treatment of the crop with 1 pound of insecticide per acre (Miles et al., 1964). A total diet study conducted in 1969-1970 revealed methoxychlor concentrations ranging from 0.008 to 0.05 ppm (Corneliussen, 1972) while total U.S. dietary intake in 1970 amounted to 0.02 µg/kg body weight/day (Duggan and Corneliussen, 1972).

Exposure of the general population through ingestion of contaminated beef or milk probably contributes minimally to total dietary intake since methoxychlor undergoes rapid biodegradation in mammals and shows little tendency to bioaccumulate in tissues (International Agency for Research on Cancer, 1974). An average level of 0.001 ppm methoxychlor was detected in dairy products in the 1960s (Duggan, 1967) while no residues were detected in the body fat of cattle fed 25 ppm methoxychlor in the diet for 16 weeks (Claborn, 1956).

Methoxychlor is generally considered a relatively safe pesticide with a low order of toxicity to humans and other warm blooded animals (Farm Chemicals Handbook, 1976). However, available chronic toxicity studies for assessing carcinogenic potential are either limited in scope or inconclusive. Evaluation of carcinogenicity following skin application or subcutaneous injection was performed only in the mouse.

No tumors were observed in C3H/Anf mice following weekly skin application of 0.1 or 10 mg of methoxychlor in acetone for a total of up to 980 mg or following a single subcutaneous injection of 10 mg of the pesticide in trioctanoin (Hodge et al., 1966).

II. MATERIALS AND METHODS

A. Chemicals

Technical-grade methoxychlor [1,1'-(2,2,2-trichloroethylidene)-bis(4-methoxy)-benzene] was purchased and analyzed by Hazleton Laboratories America, Inc., Vienna, Virginia. The observed melting point (86.5° to 87.5°C) suggested a compound of high purity. Gas-liquid chromatography suggested a purity greater than 95 percent but also indicated the presence of four minor contaminants. Analyses performed 1 and 2 years later showed no substantial change in the results and thus suggested that no decomposition had occurred.

Throughout this report, the term methoxychlor is used to represent this technical-grade material.

B. Dietary Preparation

The basal laboratory diet for both control and treated animals consisted of 2 percent Duke's[®] corn oil (S.F. Sauer Company) by weight added to Wayne Lab-Blox[®] meal (Allied Mills, Inc.). Fresh mixtures of methoxychlor in corn oil were prepared weekly and stored in the dark. The mixtures of methoxychlor in corn oil were incorporated into the appropriate amount of basal laboratory diet in a twin-shell blender fitted with an accelerator bar.

C. Animals

Two animal species, rats and mice, were used in the carcinogenicity bioassay. The Osborne-Mendel rat was selected on the basis of

a comparative study of the tumorigenic responsiveness to carbon tetrachloride of five different strains of rats (Reuber and Glover, 1970). The B6C3F1 mouse was selected because it has been used by the NCI for carcinogenesis bioassays and has proved satisfactory in this capacity.

Rats and mice of both sexes were obtained through contracts of the Division of Cancer Treatment, National Cancer Institute. The Osborne-Mendel rats were procured from the Battelle Memorial Institute, Columbus, Ohio, and the B6C3F1 mice were obtained from the Charles River Breeding Laboratories, Inc., Wilmington, Massachusetts. Upon receipt, animals were quarantined for at least 10 days, observed for visible signs of disease or parasites, and assigned to the various treatment and control groups.

D. Animal Maintenance

All animals were housed by species in temperature- and humidity-controlled rooms. The temperature range was 20° to 24°C and the relative humidity was maintained between 45 and 55 percent. The air conditioning system in the laboratory provided filtered air at a rate of 12 to 15 complete changes of room air per hour. Fluorescent lighting was provided on 12-hour-daily cycle. The rats were individually housed in suspended galvanized-steel wire-mesh cages with perforated floors. Mice were housed by sex in groups of 10 in solid-bottom polypropylene cages equipped with filter tops. Sanitized cages with fresh bedding (Sanichips[®], Shurfire) were provided once each week for mice. Rats received sanitized cages with no

bedding with the same frequency. Food hoppers were changed and heat-sterilized once a week for the first 10 weeks and once a month thereafter. Fresh heat-sterilized glass water bottles were provided three times a week. Food and water were available ad libitum.

The methoxychlor-treated and control rats were housed in the same room with other rats receiving diets treated with^{*} p,p'-DDE (72-55-9) and safrole (94-59-7). All mice used in the methoxychlor study, including controls, were housed in the same room as other mice receiving diets treated with trifluralin (1582-09-8), dioxathion (78-34-2), sulfallate (95-06-7), p,p'-DDT (50-29-3), chlorobenzilate (510-15-6), p,p'-DDE (72-55-9), p,p'-TDE (72-54-8), dicofol (115-32-2), pentachloronitrobenzene (82-68-8), amitrole (61-82-5), acetaminofluorene (53-96-3), clonitralid (1420-04-8), nitrofen (1836-75-5), endosulfan (115-29-7), mexacarbate (315-18-4), and safrole (94-59-7).

E. Selection of Initial Concentrations

In order to establish the maximum tolerated concentrations of methoxychlor for administration to treated animals in the chronic studies, subchronic toxicity tests were conducted with both rats and mice. Animals of each species were distributed among six groups, each consisting of five males and five females. Methoxychlor was premixed with a small amount of corn oil. This mixture was then incorporated into the laboratory diet and fed ad libitum to five of the six rat groups at concentrations of 316, 562, 1000, 1780, and

* CAS registry numbers are given in parentheses.

3160 ppm and to five of the six mouse groups at concentrations of 1000, 1780, 3160, 5620, and 10,000 ppm. The sixth group of each species served as a control group, receiving only the basal laboratory diet. The dosed dietary preparations were administered for a period of 6 weeks, followed by a 2-week observation period during which all animals were fed the basal diet of corn oil and laboratory chow.

A dosage inducing no mortality and resulting in a depression in mean group body weight of approximately 20 percent relative to controls was to be selected as the initial high concentration. When weight gain criteria were not applicable, mortality data alone were utilized.

The initial high concentrations selected for the chronic study for male and female rats were 720 and 1500 ppm, respectively. The initial high concentrations selected for the chronic study for mice were 2800 ppm for males and 1500 ppm for females.

F. Experimental Design

The experimental design parameters for the chronic study (species, sex, group size, concentrations administered, duration of treated and untreated observation periods, and the time-weighted average concentrations) are summarized in Tables 1 and 2.

The high dose, low dose, and control rats were all approximately 6 weeks old at the time they were placed on test. The high and low concentrations of methoxychlor initially utilized for males were 720

TABLE 1
DESIGN SUMMARY FOR OSBORNE-MENDEL RATS
METHOXYCHLOR FEEDING EXPERIMENT

	INITIAL GROUP SIZE	METHOXYCHLOR CONCENTRATION ^a	OBSERVATION PERIOD		TIME-WEIGHTED AVERAGE CONCENTRATION ^b
			TREATED (WEEKS)	UNTREATED (WEEKS)	
<u>MALE</u>					
CONTROL	20	0		111	0
LOW DOSE	50	360 500 0	29 49	33	448
HIGH DOSE	50	720 1000 1000 ^c 0	29 29 16	4 33	845
<u>FEMALE</u>					
CONTROL	20	0		111	0
LOW DOSE	50	750 0	78	33	750
HIGH DOSE	50	1500 1500 ^c 0	55 17	6 34	1385

^aConcentrations given in parts per million.

^bTime-weighted average concentration = $\frac{\sum(\text{concentration} \times \text{weeks received})}{\sum(\text{weeks receiving treatment})}$

^cThese concentrations were cyclically administered with a pattern of 1 dosage-free week followed by 4 weeks of treatment at the level indicated.

TABLE 2

DESIGN SUMMARY FOR B6C3F1 MICE
METHOXYCHLOR FEEDING EXPERIMENT

	INITIAL GROUP SIZE	METHOXYCHLOR CONCENTRATION ^a	OBSERVATION PERIOD		TIME-WEIGHTED AVERAGE CONCENTRATION ^b
			TREATED (WEEKS)	UNTREATED (WEEKS)	
<u>MALE</u>					
CONTROL	20	0		92	
<hr/>					
LOW DOSE	50	1400	1		1746
		1750	77		
		0		14	
<hr/>					
HIGH DOSE	50	2800	1		3491
		3500	77		
		0		15	
<hr/>					
<hr/>					
<u>FEMALE</u>					
CONTROL	20	0		92	0
<hr/>					
LOW DOSE	50	750	1		997
		1000	77		
		0		15	
<hr/>					
HIGH DOSE	50	1500	1		1994
		2000	77		
		0		15	

^a Concentrations given in parts per million.

^b Time-weighted average concentration = $\frac{\sum(\text{concentration} \times \text{weeks received})}{\sum(\text{weeks receiving treatment})}$

and 360 ppm, respectively. At the end of week 30, the high concentration for males was increased to 1000 ppm and the low concentration was increased to 500 ppm. For females, the initial high and low concentrations of methoxychlor were 1500 and 750 ppm, respectively. In week 56 of the study, administration of methoxychlor to the high dose female rats ceased for 1 week and was then followed by 4 weeks of feeding at the previous concentration of 1500 ppm. This pattern of cyclic administration continued for the remainder of the treatment period. This same method of total intake reduction was employed for the high dose male rats beginning with week 59. Final observations on all rats were made 112 weeks after the experiment was initiated.

The high dose, low dose, and control mice were all approximately 6 weeks old at the time the experiment began. The high and low concentrations initially administered to male mice were 2800 and 1400 ppm, respectively. Female mice received initial high and low concentrations of 1500 and 750 ppm, respectively. At the end of week 2, the high and low concentrations of methoxychlor were increased, respectively, to 3500 and 1750 ppm for male mice and to 2000 and 1000 ppm for female mice. These dosages were maintained until the termination of treatment (week 78). Final observations on all mice were made 93 weeks after the experiment was initiated.

G. Clinical and Histopathologic Examinations

Animals were weighed immediately prior to initiation of the experiment. From the first day, all animals were inspected daily

for mortality. Body weights, food consumption, and data concerning appearance, behavior, signs of toxic effects, and incidence, size, and location of tissue masses were recorded at weekly intervals for the first 10 weeks and at monthly intervals thereafter. The presence of tissue masses was determined by observation and palpation of each animal.

During the course of this bioassay several pathology protocols were in effect, each for different periods of time. The minimum protocol required that, if possible, certain tissues were to be taken and examined histopathologically from all control animals, from any animal in which a tumor was observed during gross examination, and from at least 10 grossly normal males and 10 grossly normal females from each treated group. In addition, any tissues showing gross abnormalities were to be taken and examined histopathologically. Under later protocols, some tissues were taken from additional dosed animals. The number of animals in each group from which a tissue was examined is indicated in Appendices A through D.

A necropsy was performed on each animal regardless of whether it died, was killed when moribund, or was sacrificed at the end of the bioassay. The animals were euthanized by exsanguination under sodium pentobarbital anesthesia, and were immediately necropsied. The histopathologic examination consisted of gross and microscopic examination of major tissues, organs, or gross lesions taken from sacrificed animals and, whenever possible, from animals found dead.

Slides were prepared from the following tissues from selected animals: skin, subcutaneous tissue, lungs and bronchi, trachea, bone marrow, spleen, lymph nodes, thymus, heart, salivary gland, liver, gallbladder and bile duct (mice), pancreas, esophagus, stomach, small intestine, large intestine, kidney, urinary bladder, pituitary, adrenal, thyroid, parathyroid, pancreatic islets, testis, prostate, brain, uterus, mammary gland, and ovary.

Tissues for which slides were prepared were preserved in 10 percent buffered formalin, embedded in paraffin, sectioned, and stained with hematoxylin and eosin prior to microscopic examination. An occasional section was subjected to special staining techniques for more definitive diagnosis.

A few tissues were not examined for some animals, particularly for those that died early. Also, some animals were missing, cannibalized, or judged to be in such an advanced state of autolysis as to preclude histopathologic interpretation. Thus, the number of animals for which particular organs, tissues, or lesions were examined microscopically varies and does not necessarily represent the number of animals that were placed on experiment in each group.

H. Data Recording and Statistical Analyses

Pertinent data on this experiment have been recorded in an automatic data processing system, the Carcinogenesis Bioassay Data System (Linhart et al., 1974). The data elements include descriptive information on the chemicals, animals, experimental design, clinical

observations, survival, body weight, and individual pathologic results, as recommended by the International Union Against Cancer (Berenblum, 1969). Data tables were generated for verification of data transcription and for statistical review.

These data were analyzed using the statistical techniques described in this section. Those analyses of the experimental results that bear on the possibility of carcinogenicity are discussed in the statistical narrative sections.

Probabilities of survival were estimated by the product-limit procedure of Kaplan and Meier (1958) and are presented in this report in the form of graphs. Animals were statistically censored as of the time that they died of other than natural causes or were found to be missing; animals dying from natural causes were not statistically censored. Statistical analyses for a possible dose-related effect on survival used the method of Cox (1972) for testing two groups for equality and used Tarone's (1975) extensions of Cox's methods for testing a dose-related trend. One-tailed P-values have been reported for all tests except the departure from linearity test, which is only reported when its two-tailed P-value is less than 0.05.

The incidence of neoplastic or nonneoplastic lesions has been given as the ratio of the number of animals bearing such lesions at a specific anatomic site (numerator) to the number of animals in which that site was examined (denominator). In most instances, the denominators included only those animals for which that site was examined

histologically. However, when macroscopic examination was required to detect lesions prior to histologic sampling (e.g., skin or mammary tumors), or when lesions could have appeared at multiple sites (e.g., lymphomas), the denominators consist of the numbers of animals necropsied.

The purpose of the statistical analyses of tumor incidence is to determine whether animals receiving the test chemical developed a significantly higher proportion of tumors than did the control animals. As a part of these analyses, the one-tailed Fisher exact test (Cox, 1970, pp. 48-52) was used to compare the tumor incidence of a control group to that of a group of treated animals at each dose level. When results for a number of treated groups, k , are compared simultaneously with those for a control group, a correction to ensure an overall significance level of 0.05 may be made. The Bonferroni inequality (Miller, 1966, pp. 6-10) requires that the P -value for any comparison be less than or equal to $0.05/k$. In cases where this correction was used, it is discussed in the narrative section. It is not, however, presented in the tables, where the Fisher exact P -values are shown.

The Cochran-Armitage test for linear trend in proportions, with continuity correction (Armitage, 1971, pp. 362-365), was also used. Under the assumption of a linear trend, this test determined if the slope of the dose-response curve is different from zero at the one-tailed 0.05 level of significance. Unless otherwise noted, the direction of the significant trend was a positive dose relationship.

This method also provides a two-tailed test of departure from linear trend.

A time-adjusted analysis was applied when numerous early deaths resulted from causes that were not associated with the formation of tumors. In this analysis, deaths that occurred before the first tumor was observed were excluded by basing the statistical tests on animals that survived at least 52 weeks, unless a tumor was found at the anatomic site of interest before week 52. When such an early tumor was found, comparisons were based exclusively on animals that survived at least as long as the animal in which the first tumor was found. Once this reduced set of data was obtained, the standard procedures for analyses of the incidence of tumors (Fisher exact tests, Cochran-Armitage tests, etc.) were followed.

When appropriate, life-table methods were used to analyze the incidence of tumors. Curves of the proportions surviving without an observed tumor were computed as in Saffiotti et al. (1972). The week during which animals died naturally or were sacrificed was entered as the time point of tumor observation. Cox's methods of comparing these curves were used for two groups; Tarone's extension to testing for linear trend was used for three groups. The statistical tests for the incidence of tumors which used life-table methods were one-tailed and, unless otherwise noted, in the direction of a positive dose relationship. Significant departures from linearity ($P < 0.05$, two-tailed test) were also noted.

The approximate 95 percent confidence interval for the relative risk of each dosed group compared to its control was calculated from the exact interval on the odds ratio (Gart, 1971). The relative risk is defined as p_t/p_c where p_t is the true binomial probability of the incidence of a specific type of tumor in a treated group of animals and p_c is the true probability of the spontaneous incidence of the same type of tumor in a control group. The hypothesis of equality between the true proportion of a specific tumor in a treated group and the proportion in a control group corresponds to a relative risk of unity. Values in excess of unity represent the condition of a larger proportion in the treated group than in the control.

The lower and upper limits of the confidence interval of the relative risk have been included in the tables of statistical analyses. The interpretation of the limits is that in approximately 95 percent of a large number of identical experiments, the true ratio of the risk in a treated group of animals to that in a control group would be within the interval calculated from the experiment. When the lower limit of the confidence interval is greater than one, it can be inferred that a statistically significant result (a $P < 0.025$ one-tailed test when the control incidence is not zero, $P < 0.050$ when the control incidence is zero) has occurred. When the lower limit is less than unity but the upper limit is greater than unity, the lower limit indicates the absence of a significant result while the upper limit indicates that there is a theoretical possibility

of the induction of tumors by the test chemical which could not be detected under the conditions of this test.

III. CHRONIC TESTING RESULTS: RATS

A. Body Weights and Clinical Observations

Distinct, dose-related mean body weight depression was apparent throughout the treatment period in both males and females (Figure 1).

During the first 30 weeks of the study, the appearance and behavior of the treated rats were generally comparable with that of the controls, except that occasional hunched appearance, reddened eyes, and abdominal urine stains were noted in one to three treated rats as early as week 4.

From week 34 to cessation of treatment (week 78), some clinical signs were noted with slightly greater frequency in the treated groups than in the controls. During the post-treatment observation period these signs were observed at comparable rates in treated and control animals. These signs included hunched appearance; urine staining; body sores; rough fur; eyes that were cloudy, squinted, or reddened with discharge or crust; and palpable nodules or tissue masses. As the animals aged, the incidence of these signs in all groups gradually increased. Respiratory signs characterized by labored respiration, wheezing, and/or nasal discharge were observed at a low incidence in all groups throughout the study. Isolated, sporadic, and spontaneous observations in one to three rats included circling, small testes, head tilt, "stiff" gait, hind-limb paralysis, ataxia, and transient vaginal discharge.





FIGURE 1
GROWTH CURVES FOR METHOXYCHLOR CHRONIC STUDY RATS

B. Survival

The estimated probabilities of survival for male and female rats in the control and methoxychlor-treated groups are shown in Figure 2. For both male and female rats the Tarone test did not detect a significant positive association between increased dosage and elevated mortality.

For males the actual survival was adequate as 86 percent of the high dose, 74 percent of the low dose and 85 percent of the control rats lived at least 100 weeks. For females the actual survival was also adequate as 94 percent of the high dose, 94 percent of the low dose, and 90 percent of the control rats lived at least 100 weeks.

C. Pathology

Histopathologic findings on neoplasms in rats are tabulated in Appendix A (Tables A1 and A2); findings on nonneoplastic lesions are tabulated in Appendix C (Tables C1 and C2).

Hemangiosarcomas were the only tumors observed at unusually high incidences. The hemangiosarcomas occurred in the spleen of 1/20 control males, 6/44 low dose males, 2/42 high dose males, and 1/20 control females. Hemangiosarcomas also occurred as subcutaneous masses in 2/50 low dose males and as an abdominal tissue mass in 1/50 low dose male rats. Thus, there was an increased incidence of hemangiosarcomas at all sites in male rats receiving methoxychlor (9 low dose and 2 high dose) as compared to a single control male and a single control female with these tumors. The lack of a dose response in the





FIGURE 2
SURVIVAL COMPARISONS OF METHOXYCHLOR CHRONIC STUDY RATS

number of male rats with hemangiosarcomas makes this finding of questionable importance even though in our experience hemangiosarcomas occur relatively infrequently in the Osborne-Mendel rat. With the exception of a hemangiosarcoma in the spleen of one control female there were no unusual tumors or any unusual incidences of spontaneous tumors in the female rats.

Microscopically, the hemangiosarcomas consisted of the formation of capillaries and cavernous vascular spaces that often contained blood and were lined with hyperchromatic, plump anaplastic endothelial cells. There was piling up of these cells around the vascular spaces. These cells invaded and replaced the normal architecture of the spleen and involved tissues.

Other neoplasms observed during this examination were seen at incidences often found to occur naturally in untreated aged Osborne-Mendel rats.

Inflammatory, degenerative, and proliferative lesions as seen in the control and treated animals were similar in number and kind to those spontaneous lesions occurring in aged Osborne-Mendel rats.

There was no definitive evidence of the carcinogenicity of methoxychlor in Osborne-Mendel rats.

D. Statistical Analyses of Results

The results of the statistical analyses of tumor incidence in rats are summarized in Tables 3 and 4. The analysis for every type of tumor that was observed in more than 5 percent of any of the methoxychlor-dosed groups of either sex is included.

TABLE 3
ANALYSES OF THE INCIDENCE OF PRIMARY TUMORS AT
SPECIFIC SITES IN MALE RATS TREATED WITH METHOXYCHLOR^a

TOPOGRAPHY:MORPHOLOGY	CONTROL	LOW DOSE	HIGH DOSE
Subcutaneous Tissue: Fibroma or Fibrosarcoma ^b	1/20(0.05)	2/50(0.04)	3/50(0.06)
P Values ^c	N.S.	N.S.	N.S.
Relative Risk (Control) ^d			
Lower Limit	---	0.800	1.200
Upper Limit	---	0.045	0.105
	---	46.273	61.724
Weeks to First Observed Tumor	111	101	26
Liver: Neoplastic Nodule or Hepatocellular Carcinoma ^b	0/19(0.00)	2/37(0.05)	0/37(0.00)
P Values ^c	N.S.	N.S.	N.S.
Relative Risk (Control) ^d			
Lower Limit	---	Infinite	---
Upper Limit	---	0.157	---
	---	Infinite	---
Weeks to First Observed Tumor	---	96	---
Pancreas: Liposarcoma ^b	0/19(0.00)	1/20(0.05)	0/20(0.00)
P Values ^c	N.S.	N.S.	N.S.
Relative Risk (Control) ^d			
Lower Limit	---	Infinite	---
Upper Limit	---	0.052	---
	---	Infinite	---
Weeks to First Observed Tumor	---	79	---

TABLE 3 (Continued)

TOPOGRAPHY:MORPHOLOGY	CONTROL	LOW DOSE	HIGH DOSE
Kidney: Transitional-Cell Carcinoma ^b	0/19(0.00)	1/21(0.05)	0/28(0.00)
P Values ^c	N.S.	N.S.	N.S.
Relative Risk (Control) ^d	---	Infinite	---
Lower Limit	---	0.050	---
Upper Limit	---	Infinite	---
Weeks to First Observed Tumor	---	94	---
Kidney: Fibrous Histiocytoma, Malignant ^b	0/19(0.00)	1/21(0.05)	0/28(0.00)
P Values ^c	N.S.	N.S.	N.S.
Relative Risk (Control) ^d	---	Infinite	---
Lower Limit	---	0.050	---
Upper Limit	---	Infinite	---
Weeks to First Observed Tumor	---	96	---
Kidney: Mixed Tumor, Malignant ^b	1/19(0.05)	2/21(0.10)	1/28(0.04)
P Values ^c	N.S.	N.S.	N.S.
Relative Risk (Control) ^d	---	1.810	0.679
Lower Limit	---	0.103	0.009
Upper Limit	---	100.601	51.519
Weeks to First Observed Tumor	111	63	111

TABLE 3 (Continued)

TOPOGRAPHY: MORPHOLOGY	CONTROL	LOW DOSE	HIGH DOSE
Kidney: Hamartoma ^{*b}	0/19(0.00)	1/21(0.05)	0/28(0.00)
P Values ^c	N.S.	N.S.	N.S.
Relative Risk (Control) ^d	---	Infinitive	---
Lower Limit	---	0.050	---
Upper Limit	---	Infinitive	---
Weeks to First Observed Tumor	---	111	---
Pituitary: Chromophobe Adenoma ^b	3/18(0.17)	8/21(0.38)	4/22(0.18)
P Values ^c	N.S.	N.S.	N.S.
Relative Risk (Control) ^d	---	2.286	1.091
Lower Limit	---	0.662	0.216
Upper Limit	---	11.364	6.582
Weeks to First Observed Tumor	105	69	111
Adrenal: Pheochromocytoma ^b	0/20(0.00)	1/18(0.06)	1/19(0.05)
P Values ^c	N.S.	N.S.	N.S.
Relative Risk (Control) ^d	---	Infinitive	Infinitive
Lower Limit	---	0.061	0.058
Upper Limit	---	Infinitive	Infinitive
Weeks to First Observed Tumor	---	111	111

^{*}This is considered to be a benign form of the malignant mixed tumor of the kidney and consists of proliferative lipocytes, tubular structures, fibroblasts, and vascular spaces in varying proportions.

TABLE 3 (Continued)

TOPOGRAPHY:MORPHOLOGY	CONTROL	LOW DOSE	HIGH DOSE
Thyroid: Follicular-Cell Carcinoma ^b	0/19(0.00)	4/27(0.15)	1/24(0.04)
P Values ^c	N.S.	N.S.	N.S.
Departure from Linear Trend ^e	P = 0.044	---	---
Relative Risk (Control) ^d	---	Infinitive	Infinitive
Lower Limit	---	0.685	0.044
Upper Limit	---	Infinitive	Infinitive
Weeks to First Observed Tumor	---	69	111
Thyroid: Follicular-Cell Adenoma ^b	1/19(0.05)	1/27(0.04)	2/24(0.08)
P Values ^c	N.S.	N.S.	N.S.
Relative Risk (Control) ^d	---	0.704	1.583
Lower Limit	---	0.010	0.090
Upper Limit	---	53.206	89.314
Weeks to First Observed Tumor	105	111	111
Thyroid: Follicular-Cell Adenoma or Follicular-Cell Carcinoma ^b	1/19(0.05)	5/27(0.19)	3/24(0.13)
P Values ^c	N.S.	N.S.	N.S.
Relative Risk (Control) ^d	---	3.519	2.375
Lower Limit	---	0.447	0.212
Upper Limit	---	159.091	118.982
Weeks to First Observed Tumor	105	69	111

TABLE 3 (Continued)

TOPOGRAPHY: MORPHOLOGY	CONTROL	LOW DOSE	HIGH DOSE
Thyroid: C-Cell Adenoma ^b	2/19(0.11)	0/27(0.00)	2/24(0.08)
P Values ^c	N.S.	N.S.	N.S.
Relative Risk (Control) ^d	---	0.000	0.792
Lower Limit	---	0.000	0.063
Upper Limit	---	2.319	10.089
Weeks to First Observed Tumor	109	---	110
Thyroid: C-Cell Adenoma or C-Cell Carcinoma ^b	2/19(0.11)	1/27(0.04)	3/24(0.13)
P Values ^c	N.S.	N.S.	N.S.
Relative Risk (Control) ^d	---	0.352	1.188
Lower Limit	---	0.006	0.153
Upper Limit	---	6.315	13.079
Weeks to First Observed Tumor	109	111	110
Pancreatic Islets: Islet-Cell Adenoma ^b	0/19(0.00)	1/20(0.05)	1/20(0.05)
P Values ^c	N.S.	N.S.	N.S.
Relative Risk (Control) ^d	---	Infinite	Infinite
Lower Limit	---	0.052	0.052
Upper Limit	---	Infinite	Infinite
Weeks to First Observed Tumor	---	101	111

TABLE 3 (Concluded)

TOPOGRAPHY: MORPHOLOGY	CONTROL	LOW DOSE	HIGH DOSE
Testis: Interstitial-Cell Tumor ^b	0/20(0.00)	0/19(0.00)	1/21(0.05)
P Values ^c	N.S.	N.S.	N.S.
Relative Risk (Control) ^d	---	---	Infinite
Lower Limit	---	---	0.052
Upper Limit	---	---	Infinite
Weeks to First Observed Tumor	---	---	111
Circulatory System: Hemangiosarcoma ^b	1/20(0.05)	9/50(0.18)	2/50(0.04)
P Values ^c	N.S.	N.S.	N.S.
Departure from Linear Trend ^e	P = 0.020	---	---
Relative Risk (Control) ^d	---	3.600	0.800
Lower Limit	---	0.562	0.045
Upper Limit	---	154.106	46.273
Weeks to First Observed Tumor	108	87	111

^aTreated groups received time-weighted average doses of 448 or 845 ppm in feed.

^bNumber of tumor-bearing animals/number of animals examined at site (proportion).

^cThe probability level for the Cochran-Armitage test is given beneath the incidence of tumors in the control group when $P < 0.05$; otherwise, not significant (N.S.) is indicated. The probability level for the Fisher exact test for the comparison of a treated group with the control group is given beneath the incidence of tumors in the treated group when $P < 0.05$; otherwise, not significant (N.S.) is indicated. For both Cochran-Armitage and Fisher exact tests a negative designation (N) indicates a lower incidence in the treated group(s) than in the control group.

^dThe 95% confidence interval on the relative risk of the treated group to the control group.

^eThe probability level of the test for departure from linear trend is given beneath the control group when $P < 0.05$.

TABLE 4
ANALYSES OF THE INCIDENCE OF PRIMARY TUMORS AT
SPECIFIC SITES IN FEMALE RATS TREATED WITH METHOXYCHLOR^a

TOPOGRAPHY: MORPHOLOGY	CONTROL	LOW DOSE	HIGH DOSE
Subcutaneous Tissue: Fibroma ^b	1/20(0.05)	4/50(0.08)	0/50(0.00)
P Values ^c	N.S.	N.S.	N.S.
Relative Risk (Control) ^d	---	1.600	0.000
Lower Limit	---	0.175	0.000
Upper Limit	---	77.169	7.475
Weeks to First Observed Tumor	94	100	---
Subcutaneous Tissue: Fibroma or Fibrosarcoma ^b	1/20(0.05)	5/50(0.10)	1/50(0.02)
P Values ^c	N.S.	N.S.	N.S.
Relative Risk (Control) ^d	---	2.000	0.400
Lower Limit	---	0.249	0.005
Upper Limit	---	92.596	30.802
Weeks to First Observed Tumor	94	100	72
Pituitary: Chromophobe Adenoma ^b	8/20(0.40)	7/32(0.22)	10/36(0.28)
P Values ^c	N.S.	N.S.	N.S.
Relative Risk (Control) ^d	---	0.547	0.694
Lower Limit	---	0.209	0.308
Upper Limit	---	1.474	1.722
Weeks to First Observed Tumor	106	95	106

TABLE 4 (Continued)

TOPOGRAPHY: MORPHOLOGY	CONTROL	LOW DOSE	HIGH DOSE
Thyroid: Follicular-Cell Adenoma ^b	0/20(0.00)	2/28(0.07)	0/31(0.00)
P Values ^c	N.S.	N.S.	N.S.
Relative Risk (Control) ^d	---	Infinite	---
Lower Limit	---	0.220	---
Upper Limit	---	Infinite	---
Weeks to First Observed Tumor	---	111	---
Thyroid: C-Cell Adenoma ^b	0/20(0.00)	0/28(0.00)	3/31(0.10)
P Values ^c	N.S.	N.S.	N.S.
Relative Risk (Control) ^d	---	---	Infinite
Lower Limit	---	---	0.404
Upper Limit	---	---	Infinite
Weeks to First Observed Tumor	---	---	112
Thyroid: C-Cell Adenoma or C-Cell Carcinoma ^b	0/20(0.00)	1/28(0.04)	3/31(0.10)
P Values ^c	N.S.	N.S.	N.S.
Relative Risk (Control) ^d	---	Infinite	Infinite
Lower Limit	---	0.039	0.404
Upper Limit	---	Infinite	Infinite
Weeks to First Observed Tumor	---	111	112

TABLE 4 (Continued)

TOPOGRAPHY:MORPHOLOGY	CONTROL	LOW DOSE	HIGH DOSE
Mammary Gland: Adenocarcinoma NOS ^b	1/20(0.05)	1/50(0.02)	3/50(0.06)
P Values ^c	N.S.	N.S.	N.S.
Relative Risk (Control) ^d	---	0.400	1.200
Lower Limit	---	0.005	0.105
Upper Limit	---	30.802	61.724
Weeks to First Observed Tumor	111	16	106
Mammary Gland: Adenoma or Adenocarcinoma NOS ^b	1/20(0.05)	1/50(0.02)	5/50(0.10)
P Values ^c	N.S.	N.S.	N.S.
Relative Risk (Control) ^d	---	0.400	2.000
Lower Limit	---	0.005	0.249
Upper Limit	---	30.802	92.596
Weeks to First Observed Tumor	111	16	106
Uterus: Endometrial Stromal Polyp ^b	0/19(0.00)	1/34(0.03)	4/35(0.11)
P Values ^c	N.S.	N.S.	N.S.
Relative Risk (Control) ^d	---	Infinite	Infinite
Lower Limit	---	0.031	0.526
Upper Limit	---	Infinite	Infinite
Weeks to First Observed Tumor	---	111	111

TABLE 4 (Concluded)

- a. Treated groups received time-weighted average doses of 750 or 1385 ppm in feed.
- b. Number of tumor-bearing animals/number of animals examined at site (proportion).
- c. The probability level for the Cochran-Armitage test is given beneath the incidence of tumors in the control group when $P < 0.05$; otherwise, not significant (N.S.) is indicated. The probability level for the Fisher exact test for the comparison of a treated group with the control group is given beneath the incidence of tumors in the treated group when $P < 0.05$; otherwise, not significant (N.S.) is indicated. For both Cochran-Armitage and Fisher exact tests a negative designation (N) indicates a lower incidence in the treated group(s) than in the control group.
- d. The 95% confidence interval on the relative risk of the treated group to the control group.

None of the statistical tests for rats of either sex indicated a significant positive association between the administration of methoxychlor and an elevated tumor incidence. Thus, at the dose levels used in this experiment there was no statistical evidence that methoxychlor was a carcinogen in Osborne-Mendel rats.

To provide additional insight, 95 percent confidence intervals on the relative risk have been estimated and entered in the tables based upon the observed tumor incidence rates. In many of the intervals shown in Tables 3 and 4, the value one is included; this indicates the absence of statistically significant results. It should also be noted that many of the confidence intervals have an upper limit greater than one, indicating the theoretical possibility of a significantly increased rate of tumor incidence induced in rats by methoxychlor that could not be established under the conditions of this test.

IV. CHRONIC TESTING RESULTS: MICE

A. Body Weights and Clinical Observations

Distinct, dose-related mean body weight depression was apparent in treated females during the entire bioassay. There were no distinct differences in weight gain patterns of low dose and control male mice; high dose male mice gained slightly less weight than these groups during the treatment period (Figure 3).

Throughout the study there was no definitive evidence of compound effects with regard to physical appearance and behavior of the treated mice. Signs often observed in group-housed laboratory mice, particularly in males, were observed at a comparable rate in treated and control animals. These common signs included sores on the body and/or extremities (more prevalent in the males); a hunched appearance; localized alopecia; penile, vulvar, or anal irritation with apparent sores and abscesses around the inguinal area (more prevalent in male mice); rough or stained fur; bloating; and palpable nodules or tissue masses. Isolated observations in one or two mice from all groups included reddened or squinted eyes and labored respiration.

B. Survival

The estimated probabilities of survival for male and female mice in the control and methoxychlor-treated groups are shown in Figure 4. For both male and female mice the Tarone test did not detect a significant positive association between increased dosage and elevated mortality.





FIGURE 3
GROWTH CURVES FOR METHOXYCHLOR CHRONIC STUDY MICE





FIGURE 4
SURVIVAL COMPARISONS OF METHOXYCHLOR CHRONIC STUDY MICE

For males the actual survival was adequate as 69 percent of the high dose, 58 percent of the low dose, and 45 percent of the control mice survived over 81 weeks. For females the actual survival was high as 98 percent of the high dose, 90 percent of the low dose, and 85 percent of the control mice survived until the end of the test.

C. Pathology

Histopathologic findings on neoplasms in mice are tabulated in Appendix B (Tables B1 and B2); findings on nonneoplastic lesions are tabulated in Appendix D (Tables D1 and D2).

A variety of neoplasms were observed among both treated and control mice. Each of the types of tumors represented has been encountered previously as a naturally occurring lesion in B6C3F1 mice, and is without apparent relationship to the administration of the chemical.

The inflammatory, degenerative, and proliferative lesions that occurred in the control and treated animals were also without appreciable differences in number and kind from those spontaneous lesions occurring in aged B6C3F1 mice.

No evidence for the carcinogenicity of methoxychlor in B6C3F1 mice was provided by this histopathologic examination.

D. Statistical Analyses of Results

The results of the statistical analyses of tumor incidence in mice are summarized in Tables 5 and 6. The analysis for every type

TABLE 5

ANALYSES OF THE INCIDENCE OF PRIMARY TUMORS AT
SPECIFIC SITES IN MALE MICE TREATED WITH METHOXYCHLOR^a

TOPOGRAPHY: MORPHOLOGY	CONTROL	LOW DOSE	HIGH DOSE
Subcutaneous Tissue: Fibrosarcoma ^b	1/12 (0.08)	2/44 (0.05)	2/47 (0.04)
P Values ^c	N.S.	N.S.	N.S.
Relative Risk (Control) ^d	---	0.545	0.511
Lower Limit	---	0.032	0.031
Upper Limit	---	31.472	29.570
Weeks to First Observed Tumor	92	86	71
Liver: Hepatocellular Carcinoma ^b	3/13 (0.23)	3/36 (0.08)	6/32 (0.19)
P Values ^c	N.S.	N.S.	N.S.
Relative Risk (Control) ^d	---	0.361	0.813
Lower Limit	---	0.058	0.216
Upper Limit	---	2.452	4.490
Weeks to First Observed Tumor	91	85	92

^aTreated groups received time-weighted average doses of 1746 or 3491 ppm in feed.

^bNumber of tumor-bearing animals/number of animals examined at site (proportion).

^cThe probability level for the Cochran-Armitage test is given beneath the incidence of tumors in the control group when $P < 0.05$; otherwise, not significant (N.S.) is indicated. The probability level for the Fisher exact test for the comparison of a treated group with the control group is given beneath the incidence of tumors in the treated group when $P < 0.05$; otherwise, not significant (N.S.) is indicated. For both Cochran-Armitage and Fisher exact tests a negative designation (N) indicates a lower incidence in the treated group(s) than in the control group.

^dThe 95% confidence interval on the relative risk of the treated group to the control group.

TABLE 6
ANALYSES OF THE INCIDENCE OF PRIMARY TUMORS AT
SPECIFIC SITES IN FEMALE MICE TREATED WITH METHOXYCHLOR^a

TOPOGRAPHY:MORPHOLOGY	CONTROL	LOW DOSE	HIGH DOSE
Hematopoietic System: Malignant Lymphoma ^b	1/20(0.05)	5/46(0.11)	1/50(0.02)
P Values ^c	N.S.	N.S.	N.S.
Relative Risk (Control) ^d	---	2.174	0.400
Lower Limit	---	0.271	0.005
Upper Limit	---	100.415	30.802
Weeks to First Observed Tumor	92	79	93
Liver: Hepatocellular Carcinoma ^b	0/20(0.00)	1/17(0.06)	0/14(0.00)
P Values ^c	N.S.	N.S.	N.S.
Relative Risk (Control) ^d	---	Infinite	---
Lower Limit	---	0.064	---
Upper Limit	---	Infinite	---
Weeks to First Observed Tumor	---	93	---
Uterus: Endometrial Stromal Polyp ^b	1/20(0.05)	0/17(0.00)	1/13(0.08)
P Values ^c	N.S.	N.S.	N.S.
Relative Risk (Control) ^d	---	0.000	1.538
Lower Limit	---	0.000	0.021
Upper Limit	---	21.164	111.057
Weeks to First Observed Tumor	92	---	93

TABLE 6 (Concluded)

TOPOGRAPHY: MORPHOLOGY	CONTROL	LOW DOSE	HIGH DOSE
Ovary: Cystadenoma NOS ^b	0/20 (0.00)	0/17 (0.00)	1/14 (0.07)
P Values ^c	N.S.	N.S.	N.S.
Relative Risk (Control) ^d	---	---	Infinite
Lower Limit	---	---	0.079
Upper Limit	---	---	Infinite
Weeks to First Observed Tumor	---	---	93

^aTreated groups received time-weighted average doses of 997 or 1994 ppm in feed.

^bNumber of tumor-bearing animals/number of animals examined at site (proportion).

^cThe probability level for the Cochran-Armitage test is given beneath the incidence of tumors in the control group when $P < 0.05$; otherwise, not significant (N.S.) is indicated. The probability level for the Fisher exact test for the comparison of a treated group with the control group is given beneath the incidence of tumors in the treated group when $P < 0.05$; otherwise, not significant (N.S.) is indicated. For both Cochran-Armitage and Fisher exact tests a negative designation (N) indicates a lower incidence in the treated group(s) than in the control group.

^dThe 95% confidence interval on the relative risk of the treated group to the control group.

of tumor that was observed in more than 5 percent of any of the methoxychlor-dosed groups of either sex is included.

None of the statistical tests for mice of either sex indicated a significant positive association between the administration of methoxychlor and increased tumor incidence. Thus, at the dose levels used in this experiment, there was no statistical evidence that methoxychlor was a carcinogen in B6C3F1 mice.

To provide additional insight, 95 percent confidence intervals on the relative risk have been estimated and entered in the tables based upon the observed tumor incidence rates. In many of the intervals shown in Tables 5 and 6, the value one is included; this indicates the absence of statistically significant results. It should also be noted that many of the confidence intervals have an upper limit greater than one, indicating the theoretical possibility of a significantly increased rate of tumor incidence induced in mice by methoxychlor that could not be established under the conditions of this test.

V. DISCUSSION

Under the conditions of this bioassay, dietary administration of methoxychlor was not associated with an increased incidence of any type of neoplasm in rats or mice of either sex. No statistically significant positive association was demonstrated between dosage and mortality. In all treated groups there was dose-related mean body weight depression when compared to controls.

Although hemangiosarcomas occur relatively infrequently in Osborne-Mendel rats, the occurrence of these neoplasms in males (nine low dose and two high dose) in this bioassay is not considered to be compound-related, primarily due to the absence of a dose-related response. No significant statistical association was established between methoxychlor treatment and the incidence of any tumor in either male or female rats.

The pathologist detected no tumors of histopathologic significance in mice of either sex, and no tumors in dosed mice were observed at statistically significant incidences.

Methoxychlor has been previously assayed by the oral route in the rat. In two studies (Radomski et al., 1965; Deichmann et al., 1967), the incidence and distribution of both benign and malignant tumors were similar in treated and control groups; while in a third study (Hodge et al., 1952) the variety of tumors observed in the treated animals, but not observed in the controls (with the exception of one adenocarcinoma of the pancreas), was attributed to coincidence

since all these tumors commonly occurred in this particular colony of rats. None of these three studies presented evidence of the carcinogenicity of methoxychlor in rats.

Under the conditions of the present bioassay, methoxychlor was not found to be carcinogenic in Osborne-Mendel rats or B6C3F1 mice.

VI. BIBLIOGRAPHY

- Andrilenas, P.A., Farmers' Use of Pesticides in 1971--Quantities. Economic Research Service, U.S. Department of Agriculture, Agricultural Economic Report No. 252, Washington, D.C., 1974.
- Armitage, P., Statistical Methods in Medical Research, Chapter 14. J. Wiley & Sons, New York, 1971.
- Berenblum, I., editor, Carcinogenicity Testing. International Union Against Cancer, Technical Report Series, Vol. 2. International Union Against Cancer, Geneva, 1969.
- Brooks, G.T., Chlorinated Insecticides. Volume I: Technology and Application. CRC Press, Inc., Cleveland, Ohio, 1974.
- Chemical Abstracts Service. The Chemical Abstracts Service (CAS) Ninth Collective Index, Volumes 76-85, 1972-1976. American Chemical Society, Washington, D.C., 1977.
- Claborn, H.V., Insecticide Residues in Meat and Milk. U.S. Department of Agriculture ARS-33-25, U.S. Government Printing Office, Washington, D.C., 1956. (As cited in International Agency for Research on Cancer, 1974.)
- Corneliussen, P.E., "Pesticide Residues in Total Diet Samples (VI)." Pesticide Monitoring Journal 5:513, 1972. (As cited in International Agency for Research on Cancer, 1974.)
- Cox, D.R., Analysis of Binary Data, Chapters 4 and 5. Methuen and Co., Ltd., London, 1970.
- Cox, D.R. "Regression Models and Life-Tables." Journal of the Royal Statistical Society, Series "B" 34:187-220, 1972.
- Deichmann, W.B., M. Keplinger, F. Salo, and E. Glass, "Synergism Among Oral Carcinogens. IV: The Simultaneous Feeding of Four Tumorigens to Rats." Toxicology and Applied Pharmacology 11: 88, 1967. (As cited in International Agency for Research on Cancer, 1974.)
- Duggan, R.E., "Chlorinated Pesticide Residues in Fluid Milk and Other Dairy Products in the United States." Pesticide Monitoring Journal 1:2, 1967. (As cited in International Agency for Research on Cancer, 1974.)

Duggan, R.E., and P.E. Corneliussen, "Dietary Intake of Pesticide Chemicals in the United States (III), June 1968-April 1970." Pesticide Monitoring Journal 5:331, 1972. (As cited in International Agency for Research on Cancer, 1974.)

Farm Chemicals Handbook. Meister Publishing Company, Willoughby, Ohio, 1976.

Fowler, D.L., and J.N. Mahan, The Pesticide Review 1975. Agricultural Stabilization and Conservation Service, U.S. Department of Agriculture, Washington, D.C., 1976.

Gart, J.J., "The Comparison of Proportion: A Review of Significance Tests, Confidence Limits, and Adjustments for Stratification." International Statistical Institute Review 39:148-169, 1971.

Hodge, H.C., E.A. Maynard, and H.J. Blanchet, Jr., "Chronic Oral Toxicity Tests for Methoxychlor (2,2-di-[p-methoxyphenyl]-1,1,1-trichloroethane) in Rats and Dogs." Journal of Pharmacology and Experimental Therapeutics 104:60, 1952. (As cited in International Agency for Research on Cancer, 1974.)

Hodge, H.C., E.A. Maynard, W.L. Downs, J.K. Ashton, and L.L. Salerno, "Tests on Mice for Evaluating Carcinogenicity." Toxicology and Applied Pharmacology 9:583, 1966. (As cited in International Agency for Research on Cancer, 1974.)

International Agency for Research on Cancer, "Methoxychlor." IARC Monographs on the Evaluation of Carcinogenic Risk of Chemicals to Man. Volume 5: Some Organochlorine Pesticides. Lyon, France, 1974.

Kaplan, E.L., and P. Meier, "Nonparametric Estimation from Incomplete Observations." Journal of the American Statistical Association 53:457-481, 1958.

Linhart, M.S., J.A. Cooper, R.L. Martin, N.P. Page, and J.A. Peters, "Carcinogenesis Bioassay Data System." Computers and Biomedical Research 7:230-248, 1974.

Miles, J.R.W., W.W. Sans, H.B. Wressell, and G.F. Manson, "Growth-Dilution as a Factor in the Decline of Pesticide Residues of Alfalfa Grass Forage." Canadian Journal of Plant Science 44:37, 1964. (As cited in International Agency for Research on Cancer, 1974.)

Miller, R.G., Simultaneous Statistical Inference. McGraw-Hill Book Co., New York, 1966.

- Ouellette, R.P., and J. King, "Pesticides '76." Chemical Week 118: 27-38, June 23, 1976.
- Radomski, J.L., W.B. Deichmann, W.E. MacDonald, and E.M. Glass, "Synergism Among Oral Carcinogens. I: Results of the Simultaneous Feeding of Four Tumorigens to Rats." Toxicology and Applied Pharmacology 7:652, 1965. (As cited in International Agency for Research on Cancer, 1974.)
- Reuber, M.D., and E.L. Glover, "Cirrhosis and Carcinoma of the Liver in Male Rats Given Subcutaneous Carbon Tetrachloride." Journal of the National Cancer Institute 44:419-423, 1970.
- Saffiotti, U., R. Montesano, A.R. Sellakumar, F. Cefis, and D.G. Kaufman, "Respiratory Tract Carcinogenesis in Hamsters Induced by Different Numbers of Administration of Benzo (a) Pyrene and Ferric Oxide." Cancer Research 32:1073-1079, 1972.
- Spencer, E.Y., Guide to the Chemicals Used in Crop Protection, Sixth edition. Research Branch, Agriculture Canada, Publication 1093, 1973.
- Tarone, R.E., "Tests for Trend in Life-Table Analysis." Biometrika 62:679-682, 1975.
- U.S. International Trade Commission, Synthetic Organic Chemicals: United States Production and Sales 1975. USITC Publication 804, Washington, D.C., 1977.

APPENDIX A

SUMMARY OF THE INCIDENCE OF NEOPLASMS
IN RATS TREATED WITH METHOXYCHLOR

123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000

TABLE A1
SUMMARY OF THE INCIDENCE OF NEOPLASMS IN MALE RATS TREATED WITH METHOXYCHLOR

	CONTROL (VEH) 01-M023	LOW DOSE 01-M024	HIGH DOSE 01-M025
ANIMALS INITIALLY IN STUDY	20	50	50
ANIMALS NECROPSIED	20	50	50
ANIMALS EXAMINED HISTOPATHOLOGICALLY **	20	44	41
INTEGUMENTARY SYSTEM			
*SKIN	(20)	(50)	(50)
FIBROSARCOMA			1 (2%)
*SUBCUT TISSUE	(20)	(50)	(50)
FIBROMA	1 (5%)	2 (4%)	2 (4%)
FIBROSARCOMA			1 (2%)
LIPOSARCOMA	1 (5%)		
HEMANGIOSARCOMA		2 (4%)	
RESPIRATORY SYSTEM			
#LUNG	(20)	(20)	(23)
HEPATOCELLULAR CARCINOMA, METAST		1 (5%)	
FIBROSARCOMA, METASTATIC			1 (4%)
MIXED TUMOR, METASTATIC		1 (5%)	
OSTEOSARCOMA, METASTATIC	1 (5%)		1 (4%)
HEMATOPOIETIC SYSTEM			
*MULTIPLE ORGANS	(20)	(50)	(50)
MALIG.LYMPHOMA, HISTIOCYTIC TYPE			2 (4%)
*SUBCUT TISSUE/BACK	(20)	(50)	(50)
MALIG.LYMPHOMA, HISTIOCYTIC TYPE		1 (2%)	
#SPLEEN	(20)	(44)	(42)
HEMANGIOSARCOMA	1 (5%)	6 (14%)	2 (5%)
CIRCULATORY SYSTEM			
NONE			
# NUMBER OF ANIMALS WITH TISSUE EXAMINED MICROSCOPICALLY			
* NUMBER OF ANIMALS NECROPSIED			
**EXCLUDES PARTIALLY AUTOLYZED ANIMALS			

TABLE A1 (CONTINUED)

	CONTROL (VEH) 01-M023	LOW DOSE 01-M024	HIGH DOSE 01-M025
DIGESTIVE SYSTEM			
#LIVER	(19)	(37)	(37)
NEOPLASTIC NODULE		1 (3%)	
HEPATOCELLULAR CARCINOMA		1 (3%)	
#PANCREAS	(19)	(20)	(20)
LIPOSARCOMA		1 (5%)	
URINARY SYSTEM			
#KIDNEY	(19)	(21)	(28)
TRANSITIONAL-CELL CARCINOMA		1 (5%)	
FIBROUS HISTIOCYTOMA, MALIGNANT		1 (5%)	
MIXED TUMOR, MALIGNANT	1 (5%)	2 (10%)	1 (4%)
OSTEOSARCOMA, METASTATIC	1 (5%)		
HAMARTOMA+		1 (5%)	
ENDOCRINE SYSTEM			
#PITUITARY	(18)	(21)	(22)
CHROMOPHOBE ADENOMA	3 (17%)	8 (38%)	4 (18%)
#ADRENAL	(20)	(18)	(19)
CORTICAL CARCINOMA	1 (5%)		
PHEOCHROMOCYTOMA		1 (6%)	1 (5%)
#THYROID	(19)	(27)	(24)
FOLLICULAR-CELL ADENOMA	1 (5%)	1 (4%)	2 (8%)
FOLLICULAR-CELL CARCINOMA		4 (15%)	1 (4%)
C-CELL ADENOMA	2 (11%)		2 (8%)
C-CELL CARCINOMA		1 (4%)	1 (4%)
#PANCREATIC ISLETS	(19)	(20)	(20)
ISLET-CELL ADENOMA		1 (5%)	1 (5%)
REPRODUCTIVE SYSTEM			
*MAMMARY GLAND	(20)	(50)	(50)
ADENOMA, NOS		1 (2%)	1 (2%)
#TESTIS	(20)	(19)	(21)
INTERSTITIAL-CELL TUMOR			1 (5%)

NUMBER OF ANIMALS WITH TISSUE EXAMINED MICROSCOPICALLY

* NUMBER OF ANIMALS NECROPSIED

+ THIS IS CONSIDERED TO BE A BENIGN FORM OF THE MALIGNANT MIXED TUMOR OF THE KIDNEY AND CONSISTS OF PROLIFERATIVE LIPOCYTES, TUBULAR STRUCTURES, FIBROBLASTS, AND VASCULAR SPACES IN VARYING PROPORTIONS.

TABLE A 1 (CONTINUED)

	CONTROL (VEH) 01-M023	LOW DOSE 01-M024	HIGH DOSE 01-M025
*EPIDIDYMIS FIBROSARCOMA	(20)	(50)	(50) 1 (2%)
NERVOUS SYSTEM			
NONE			
SPECIAL SENSE ORGANS			
NONE			
MUSCULOSKELETAL SYSTEM			
*VERTEBRAL COLUMN OSTEOSARCOMA	(20) 2 (10%)	(50)	(50)
*BONE/UPPER EXTREMITY OSTEOSARCOMA	(20)	(50)	(50) 1 (2%)
BODY CAVITIES			
*ABDOMINAL CAVITY FIBROSARCOMA HEMANGIOSARCOMA	(20)	(50) 1 (2%)	(50) 1 (2%)
ALL OTHER SYSTEMS			
NONE			
ANIMAL DISPOSITION SUMMARY			
ANIMALS INITIALLY IN STUDY	20	50	50
NATURAL DEATH	11	17	13
MORIBUND SACRIFICE	1	1	
SCHEDULED SACRIFICE			
ACCIDENTALLY KILLED			
TERMINAL SACRIFICE	8	32	37
ANIMAL MISSING			
@ INCLUDES AUTOLYZED ANIMALS			
# NUMBER OF ANIMALS WITH TISSUE EXAMINED MICROSCOPICALLY			
* NUMBER OF ANIMALS NECROPSIED			

TABLE A1 (CONCLUDED)

	CONTROL (VEH) 01-M023	LOW DOSE 01-M024	HIGH DOSE 01-M025
TUMOR SUMMARY			
TOTAL ANIMALS WITH PRIMARY TUMORS*	11	23	21
TOTAL PRIMARY TUMORS	13	37	26
TOTAL ANIMALS WITH BENIGN TUMORS	7	13	11
TOTAL BENIGN TUMORS	7	15	14
TOTAL ANIMALS WITH MALIGNANT TUMORS	6	17	11
TOTAL MALIGNANT TUMORS	6	21	12
TOTAL ANIMALS WITH SECONDARY TUMORS#	1	2	2
TOTAL SECONDARY TUMORS	2	2	2
TOTAL ANIMALS WITH TUMORS UNCERTAIN- BENIGN OR MALIGNANT		1	
TOTAL UNCERTAIN TUMORS		1	
TOTAL ANIMALS WITH TUMORS UNCERTAIN- PRIMARY OR METASTATIC			
TOTAL UNCERTAIN TUMORS			
* PRIMARY TUMORS: ALL TUMORS EXCEPT SECONDARY TUMORS			
# SECONDARY TUMORS: METASTATIC TUMORS OR TUMORS INVASIVE INTO AN ADJACENT ORGAN			

TABLE A2
SUMMARY OF THE INCIDENCE OF NEOPLASMS IN FEMALE RATS TREATED WITH METHOXYCHLOR

	CONTROL (VEH) 01-F023	LOW DOSE 01-F026	HIGH DOSE 01-F027
ANIMALS INITIALLY IN STUDY	20	50	50
ANIMALS NECROPSIED	20	50	50
ANIMALS EXAMINED HISTOPATHOLOGICALLY**	20	47	49
INTEGUMENTARY SYSTEM			
*SUBCUT TISSUE	(20)	(50)	(50)
SQUAMOUS CELL CARCINOMA		1 (2%)	
FIBROMA	1 (5%)	4 (8%)	
FIBROSARCOMA		1 (2%)	1 (2%)
LIPOMA		1 (2%)	
RESPIRATORY SYSTEM			
#LUNG	(20)	(33)	(33)
SQUAMOUS CELL CARCINOMA, METASTA		1 (3%)	
HEPATOCELLULAR CARCINOMA, METAST	1 (5%)		
HEMATOPOIETIC SYSTEM			
*MULTIPLE ORGANS	(20)	(50)	(50)
MALIG. LYMPHOMA, LYMPHOCYTIC TYPE	1 (5%)		
#SPLEEN	(20)	(46)	(48)
HEMANGIOSARCOMA	1 (5%)		
CIRCULATORY SYSTEM			
#ENDOCARDIUM	(20)	(32)	(30)
SARCOMA, NOS		1 (3%)	
DIGESTIVE SYSTEM			
#LIVER	(20)	(46)	(47)
NEOPLASTIC NODULE		1 (2%)	1 (2%)
HEPATOCELLULAR CARCINOMA	1 (5%)		1 (2%)
# NUMBER OF ANIMALS WITH TISSUE EXAMINED MICROSCOPICALLY			
* NUMBER OF ANIMALS NECROPSIED			
**EXCLUDES PARTIALLY AUTOLYZED ANIMALS			

TABLE A2 (CONTINUED)

	CONTROL (VEH) 01-F023	LOW DOSE 01-F026	HIGH DOSE 01-FC27
*BILE DUCT BILE DUCT ADENOMA	(20)	(50)	(50) 1 (2%)
URINARY SYSTEM			
#KIDNEY HAMARTOMA +	(20) 1 (5%)	(32)	(30)
ENDOCRINE SYSTEM			
#PITUITARY CHROMOPHOBE ADENOMA	(20) 8 (40%)	(32) 7 (22%)	(36) 10 (28%)
#THYROID FOLLICULAR-CELL ADENOMA C-CELL ADENOMA C-CELL CARCINOMA	(20)	(28) 2 (7%) 1 (4%)	(31) 3 (10%)
REPRODUCTIVE SYSTEM			
*MAMMARY GLAND ADENOMA, NOS ADENOCARCINOMA, NOS FIBROADENOMA	(20) 1 (5%) 8 (40%)	(50) 1 (2%) 16 (32%)	(50) 2 (4%) 3 (6%) 9 (18%)
*VAGINA LEIOMYOSARCOMA	(20)	(50) 1 (2%)	(50)
#UTERUS ENDOMETRIAL STROMAL POLYP	(19)	(34) 1 (3%)	(35) 4 (11%)
#OVARY CYSTADENOMA, NOS FIBROSARCOMA	(19)	(30) 1 (3%)	(30) 1 (3%)
NERVOUS SYSTEM			
#CEREBRUM ASTROCYTOMA	(20)	(32)	(30) 1 (3%)
SPECIAL SENSE ORGANS			
NONE			

NUMBER OF ANIMALS WITH TISSUE EXAMINED MICROSCOPICALLY

* NUMBER OF ANIMALS NECROPSIED

+ THIS IS CONSIDERED TO BE A BENIGN FORM OF THE MALIGNANT MIXED TUMOR OF THE KIDNEY AND CONSISTS OF PROLIFERATIVE LIPOCYTES, TUBULAR STRUCTURES, FIBROBLASTS, AND VASCULAR SPACES IN VARYING PROPORTIONS.

TABLE A2 (CONTINUED)

	CONTROL (VEH) 01-F023	LOW DOSE 01-F026	HIGH DOSE 01-F027
MUSCULOSKELETAL SYSTEM			
*VERTEBRAL COLUMN OSTEOSARCCMA	(20)	(50) 1 (2%)	(50)
BODY CAVITIES			
*ABDOMINAL CAVITY FIBROMA OSTEOSARCOMA, METASTATIC	(20)	(50) 1 (2%) 1 (2%)	(50)
ALL OTHER SYSTEMS			
NONE			
ANIMAL DISPOSITION SUMMARY			
ANIMALS INITIALLY IN STUDY	20	50	50
NATURAL DEATH@	6	3	4
MORIBUND SACRIFICE	1	4	
SCHEDULED SACRIFICE			
ACCIDENTALLY KILLED			
TERMINAL SACRIFICE	13	43	46
ANIMAL MISSING			
<u>@ INCLUDES AUTOLYZED ANIMALS</u>			
# NUMBER OF ANIMALS WITH TISSUE EXAMINED MICROSCOPICALLY			
* NUMBER OF ANIMALS NECROPSIED			

TABLE A2 (CONCLUDED)

	CONTROL (VEH) 01-F023	LOW DCSE 01-F026	HIGH DOSE 01-F027
TUMOR SUMMARY			
TOTAL ANIMALS WITH PRIMARY TUMORS*	12	30	30
TOTAL PRIMARY TUMORS	22	41	37
TOTAL ANIMALS WITH BENIGN TUMORS	12	24	26
TOTAL BENIGN TUMORS	18	32	30
TOTAL ANIMALS WITH MALIGNANT TUMORS	4	8	6
TOTAL MALIGNANT TUMORS	4	8	6
TOTAL ANIMALS WITH SECONDARY TUMORS#	1	2	
TOTAL SECONDARY TUMORS	1	2	
TOTAL ANIMALS WITH TUMORS UNCERTAIN- BENIGN OR MALIGNANT		1	1
TOTAL UNCERTAIN TUMORS		1	1
TOTAL ANIMALS WITH TUMORS UNCERTAIN- PRIMARY OR METASTATIC			
TOTAL UNCERTAIN TUMORS			
* PRIMARY TUMORS: ALL TUMORS EXCEPT SECONDARY TUMORS			
# SECONDARY TUMORS: METASTATIC TUMORS OR TUMORS INVASIVE INTO AN ADJACENT ORGAN			

APPENDIX B

SUMMARY OF THE INCIDENCE OF NEOPLASMS
IN MICE TREATED WITH METHOXYCHLOR

1113

TABLE B1
SUMMARY OF THE INCIDENCE OF NEOPLASMS IN MALE MICE TREATED WITH METHOXYCHLOR

	CONTROL (VEH) 02-M024	LOW DOSE 02-M025	HIGH DOSE 02-M026
ANIMALS INITIALLY IN STUDY	20	50	50
ANIMALS MISSING			1
ANIMALS NECROPSIED	12	44	47
ANIMALS EXAMINED HISTOPATHOLOGICALLY**	12	35	31
INTEGUMENTARY SYSTEM			
*SUBCUT TISSUE	(12)	(44)	(47)
FIBROSARCOMA	1 (8%)	2 (5%)	2 (4%)
RESPIRATORY SYSTEM			
NONE			
HEMATOPCETIC SYSTEM			
*MULTIPLE ORGANS	(12)	(44)	(47)
GRANULOCYTIC LEUKEMIA			2 (4%)
#CERVICAL LYMPH NODE	(13)	(36)	(31)
FIBROSARCOMA, METASTATIC			1 (3%)
#MESENTERIC L. NODE	(13)	(36)	(31)
FIBROSARCOMA, METASTATIC			1 (3%)
CIRCULATORY SYSTEM			
NONE			
DIGESTIVE SYSTEM			
#LIVER	(13)	(36)	(32)
HEPATOCELLULAR CARCINOMA	3 (23%)	3 (8%)	6 (19%)
URINARY SYSTEM			
NONE			
# NUMBER OF ANIMALS WITH TISSUE EXAMINED MICROSCOPICALLY			
* NUMBER OF ANIMALS NECROPSIED			
**EXCLUDES PARTIALLY AUTOLYZED ANIMALS			

TABLE B1 (CONTINUED)

	CONTROL (VEH) 02-M024	LOW DOSE 02-M025	HIGH DOSE 02-M026
ENDOCRINE SYSTEM			
NONE			
REPRODUCTIVE SYSTEM			
NONE			
NERVOUS SYSTEM			
NONE			
SPECIAL SENSE ORGANS			
NONE			
MUSCULOSKELETAL SYSTEM			
NONE			
BODY CAVITIES			
NONE			
ALL OTHER SYSTEMS			
NONE			
ANIMAL DISPOSITION SUMMARY			
ANIMALS INITIALLY IN STUDY	20	50	50
NATURAL DEATH@	12	28	17
MORIBUND SACRIFICE		2	
SCHEDULED SACRIFICE			
ACCIDENTALLY KILLED			
TERMINAL SACRIFICE	8	20	32
ANIMAL MISSING			1
@ INCLUDES AUTOLYZED ANIMALS			
# NUMBER OF ANIMALS WITH TISSUE EXAMINED MICROSCOPICALLY			
* NUMBER OF ANIMALS NECROPSIED			

TABLE B1 (CONCLUDED)

	CONTROL (VEH) 02-M024	LOW DOSE 02-M025	HIGH DOSE 02-M026
TUMOR SUMMARY			
TOTAL ANIMALS WITH PRIMARY TUMORS*	3	5	9
TOTAL PRIMARY TUMORS	4	5	10
TOTAL ANIMALS WITH BENIGN TUMORS			
TOTAL BENIGN TUMORS			
TOTAL ANIMALS WITH MALIGNANT TUMORS	3	5	9
TOTAL MALIGNANT TUMORS	4	5	10
TOTAL ANIMALS WITH SECONDARY TUMORS#			1
TOTAL SECONDARY TUMORS			2
TOTAL ANIMALS WITH TUMORS UNCERTAIN- BENIGN OR MALIGNANT			
TOTAL UNCERTAIN TUMORS			
TOTAL ANIMALS WITH TUMORS UNCERTAIN- PRIMARY OR METASTATIC			
TOTAL UNCERTAIN TUMORS			
* PRIMARY TUMORS: ALL TUMORS EXCEPT SECONDARY TUMORS			
# SECONDARY TUMORS: METASTATIC TUMORS OR TUMORS INVASIVE INTO AN ADJACENT ORGAN			

TABLE B2
SUMMARY OF THE INCIDENCE OF NEOPLASMS IN FEMALE MICE TREATED WITH METHOXYCHLOR

	CONTROL (VEH) 02-F024	LOW DOSE 02-F027	HIGH DOSE 02-F028
ANIMALS INITIALLY IN STUDY	20	50	50
ANIMALS MISSING		1	
ANIMALS NECROPSIED	20	46	50
ANIMALS EXAMINED HISTOPATHOLOGICALLY**	20	16	14
INTEGUMENTARY SYSTEM			
NONE			
RESPIRATORY SYSTEM			
NONE			
HEMATOPOIETIC SYSTEM			
*MULTIPLE ORGANS	(20)	(46)	(50)
MALIG.LYMPHOMA, LYMPHOCYTIC TYPE	1 (5%)		
MALIG.LYMPHOMA, HISTIOCYTIC TYPE		4 (9%)	1 (2%)
#COLON	(20)	(17)	(14)
MALIG.LYMPHOMA, HISTIOCYTIC TYPE		1 (6%)	
CIRCULATORY SYSTEM			
NONE			
DIGESTIVE SYSTEM			
#LIVER	(20)	(17)	(14)
HEPATOCELLULAR CARCINOMA		1 (6%)	
URINARY SYSTEM			
NONE			
ENDOCRINE SYSTEM			
#ADRENAL	(19)	(17)	(13)
CORTICAL ADENOMA	1 (5%)		
# NUMBER OF ANIMALS WITH TISSUE EXAMINED MICROSCOPICALLY			
* NUMBER OF ANIMALS NECROPSIED			
**EXCLUDES PARTIALLY AUTOLYZED ANIMALS			

TABLE B2 (CONTINUED)

	CONTROL (VEH) 02-F024	LOW DOSE 02-F027	HIGH DOSE 02-F028
REPRODUCTIVE SYSTEM			
#UTERUS	(20)	(17)	(13)
ENDOMETRIAL STROMAL POLYP	1 (5%)		1 (8%)
#OVARY	(20)	(17)	(14)
CYSTADENOMA, NOS			1 (7%)
NERVOUS SYSTEM			
NONE			
SPECIAL SENSE ORGANS			
NONE			
MUSCULOSKELETAL SYSTEM			
NONE			
BODY CAVITIES			
NONE			
ALL OTHER SYSTEMS			
NCNE			
ANIMAL DISPOSITION SUMMARY			
ANIMALS INITIALLY IN STUDY	20	50	50
NATURAL DEATH@	3	5	1
MORIBUND SACRIFICE			
SCHEDULED SACRIFICE			
ACCIDENTALLY KILLED			
TERMINAL SACRIFICE	17	44	49
ANIMAL MISSING		1	
@ INCLUDES AUTOLYZED ANIMALS			
# NUMBER OF ANIMALS WITH TISSUE EXAMINED MICROSCOPICALLY			
* NUMBER OF ANIMALS NECROPSIED			

TABLE B2 (CONCLUDED)

	CONTROL (VEH) 02-F024	LOW DOSE 02-F027	HIGH DOSE 02-F028
TUMOR SUMMARY			
TOTAL ANIMALS WITH PRIMARY TUMORS*	3	6	3
TOTAL PRIMARY TUMORS	3	6	3
TOTAL ANIMALS WITH BENIGN TUMORS	2		2
TOTAL BENIGN TUMORS	2		2
TOTAL ANIMALS WITH MALIGNANT TUMORS	1	6	1
TOTAL MALIGNANT TUMORS	1	6	1
TOTAL ANIMALS WITH SECONDARY TUMORS#			
TOTAL SECONDARY TUMORS			
TOTAL ANIMALS WITH TUMORS UNCERTAIN- BENIGN OR MALIGNANT			
TOTAL UNCERTAIN TUMORS			
TOTAL ANIMALS WITH TUMORS UNCERTAIN- PRIMARY OR METASTATIC			
TOTAL UNCERTAIN TUMORS			
* PRIMARY TUMORS: ALL TUMORS EXCEPT SECONDARY TUMORS			
# SECONDARY TUMORS: METASTATIC TUMORS OR TUMORS INVASIVE INTO AN ADJACENT ORGAN			

APPENDIX C

SUMMARY OF THE INCIDENCE OF NONNEOPLASTIC
LESIONS IN RATS TREATED WITH METHOXYCHLOR

TABLE C1
SUMMARY OF THE INCIDENCE OF NONNEOPLASTIC LESIONS IN MALE RATS TREATED WITH METHOXYCHLOR

	CONTROL (VEH) 01-M023	LOW DOSE 01-M024	HIGH DOSE 01-M025
ANIMALS INITIALLY IN STUDY	20	50	50
ANIMALS NECROPSIED	20	50	50
ANIMALS EXAMINED HISTOPATHOLOGICALLY**	20	44	41
INTEGUMENTARY SYSTEM			
*SKIN	(20)	(50)	(50)
EPIDERMAL INCLUSION CYST		2 (4%)	1 (2%)
INFLAMMATION, NOS	4 (20%)		
ABSCESS, NOS	1 (5%)		
*SUBCUT TISSUE	(20)	(50)	(50)
ABSCESS, NOS	2 (10%)		1 (2%)
RESPIRATORY SYSTEM			
#LUNG	(20)	(20)	(23)
PNEUMONIA, CHRONIC MURINE	3 (15%)	3 (15%)	1 (4%)
#ALVEOLAR WALL	(20)	(20)	(23)
MINERALIZATION	1 (5%)		
HEMATOPOIETIC SYSTEM			
#BONE MARROW	(20)	(18)	(20)
METAMORPHOSIS FATTY	1 (5%)		
#SPLEEN	(20)	(44)	(42)
HYPERTROPHY, NOS			1 (2%)
HEMATOPOIESIS	1 (5%)	3 (7%)	2 (5%)
CIRCULATORY SYSTEM			
#HEART	(20)	(21)	(20)
ARTERIOSCLEROSIS, NOS	1 (5%)		
CALCIUM DEPOSIT	1 (5%)		
#MYOCARDIUM	(20)	(21)	(20)
INFLAMMATION, NOS		2 (10%)	
DEGENERATION, NOS	8 (40%)	6 (29%)	6 (30%)
# NUMBER OF ANIMALS WITH TISSUE EXAMINED MICROSCOPICALLY			
* NUMBER OF ANIMALS NECROPSIED			
**EXCLUDES PARTIALLY AUTOLYZED ANIMALS			

TABLE C1 (CONTINUED)

	CONTROL (VEH) 01-M023	LOW DOSE 01-M024	HIGH DOSE 01-M025
#ENDOCARDIUM	(20)	(21)	(20)
INFLAMMATION, NOS		2 (10%)	
HYPERPLASIA, NOS	1 (5%)		1 (5%)
*AORTA	(20)	(50)	(50)
ARTERIOSCLEROSIS, NOS	1 (5%)	1 (2%)	
MEDIAL CALCIFICATION	2 (10%)		
CALCIUM DEPOSIT		1 (2%)	
*MESENTERIC ARTERY	(20)	(50)	(50)
ARTERIOSCLEROSIS, NOS	2 (10%)		
DIGESTIVE SYSTEM			
#LIVER	(19)	(37)	(37)
CYST, NOS	1 (5%)		1 (3%)
THROMBOSIS, NOS		1 (3%)	
INFLAMMATION, NOS		3 (8%)	
INFLAMMATION, GRANULOMATOUS			1 (3%)
METAMORPHOSIS FATTY	4 (21%)	4 (11%)	3 (8%)
*BILE DUCT	(20)	(50)	(50)
HYPERPLASIA, NOS	1 (5%)		1 (2%)
#PANCREAS	(19)	(20)	(20)
PERIARTERITIS	1 (5%)		
#STOMACH	(20)	(20)	(21)
INFLAMMATION, NOS		1 (5%)	
ULCER, FOCAL	1 (5%)	2 (10%)	1 (5%)
CALCIUM DEPOSIT	3 (15%)		
URINARY SYSTEM			
#KIDNEY	(19)	(21)	(28)
INFLAMMATION, SUPPURATIVE		2 (10%)	
INFLAMMATION, CHRONIC	12 (63%)	3 (14%)	9 (32%)
NEPHROPATHY, TOXIC		1 (5%)	1 (4%)
ENDOCRINE SYSTEM			
#ADRENAL	(20)	(18)	(19)
ANGIECTASIS	1 (5%)		
# NUMBER OF ANIMALS WITH TISSUE EXAMINED MICROSCOPICALLY			
* NUMBER OF ANIMALS NECROPSIED			

TABLE CI (CONTINUED)

	CONTROL (VEH) 01-M023	LOW DCSE 01-M024	HIGH DOSE 01-M025
*THYROID CYSTIC FOLLICLES HYPERPLASIA, C-CELL	(19)	(27) 1 (4%)	(24) 1 (4%)
*PARATHYROID HYPERPLASIA, NOS	(19) 2 (11%)	(27) 3 (11%)	(23) 2 (9%)
REPRODUCTIVE SYSTEM			
*MAMMARY GLAND GALACTOCELE	(20) 1 (5%)	(50) 1 (2%)	(50)
*TESTIS INFLAMMATION, NOS CALCIUM DEPOSIT ATROPHY, NOS	(20) 1 (5%) 6 (30%)	(19) 2 (11%)	(21) 1 (5%) 8 (38%)
*EPIDIDYMISS ATROPHY, NCS	(20) 3 (15%)	(50)	(50)
NERVOUS SYSTEM			
NONE			
SPECIAL SENSE ORGANS			
*EYE CATARACT	(20)	(50) 1 (2%)	(50)
*EYE/CORNEA INFLAMMATION, NOS	(20)	(50) 1 (2%)	(50)
*EYE/CILIARY BODY INFLAMMATION, NOS	(20)	(50) 1 (2%)	(50)
*EYE/IRIS INFLAMMATION, NOS	(20)	(50) 1 (2%)	(50)
*EYE/RETINA INFLAMMATION, NOS	(20)	(50) 1 (2%)	(50)
*EYE/CONJUNCTIVA INFLAMMATION, NOS	(20)	(50) 1 (2%)	(50)

* NUMBER OF ANIMALS WITH TISSUE EXAMINED MICROSCOPICALLY

* NUMBER OF ANIMALS NECROPSIED

TABLE C1 (CONCLUDED)

	CONTROL (VEH) 01-M023	LOW DOSE 01-M024	HIGH DOSE 01-M025
MUSCULOSKELETAL SYSTEM			
*MUSCLE HIP/THIGH INFLAMMATION, NOS DEGENERATION, NOS	(20)	(50) 1 (2%)	(50) 1 (2%)
BODY CAVITIES			
*ABDOMINAL CAVITY NECROSIS, FAT	(20) 1 (5%)	(50) 3 (6%)	(50) 1 (2%)
*PERICARDIUM INFLAMMATION, NOS	(20) 1 (5%)	(50) 1 (2%)	(50)
*MESENTERY PERIARTERITIS NECROSIS, FAT	(20) 1 (5%)	(50) 1 (2%)	(50) 1 (2%)
ALL OTHER SYSTEMS			
NONE			
SPECIAL MORPHOLOGY SUMMARY			
NO LESION REPORTED	1	12	10
NECROPSY PERF/NO HISTO PERFORMED		6	6
AUTO/NECROPSY/HISTO PERF		1	
AUTO/NECROPSY/NO HISTO			3
# NUMBER OF ANIMALS WITH TISSUE EXAMINED MICROSCOPICALLY			
* NUMBER OF ANIMALS NECROPSIED			

TABLE C2
SUMMARY OF THE INCIDENCE OF NONNEOPLASTIC LESIONS IN FEMALE RATS TREATED WITH METHOXYCHLOR

	CONTROL (VEH) 01-F023	LOW DCSE 01-F026	HIGH DOSE 01-F027
ANIMALS INITIALLY IN STUDY	20	50	50
ANIMALS NECROPSIED	20	50	50
ANIMALS EXAMINED HISTOPATHOLOGICALLY **	20	47	49
INTEGUMENTARY SYSTEM			
NONE			
RESPIRATORY SYSTEM			
#LUNG	(20)	(33)	(33)
INFLAMMATION, SUPPURATIVE			1 (3%)
PNEUMONIA, CHRONIC MURINE	3 (15%)	1 (3%)	2 (6%)
HEMATOPOIETIC SYSTEM			
#SPLEEN	(20)	(46)	(48)
HEMATOPOIESIS	3 (15%)	1 (2%)	2 (4%)
CIRCULATORY SYSTEM			
#MYOCARDIUM	(20)	(32)	(30)
DEGENERATION, NOS	2 (10%)	4 (13%)	4 (13%)
#ENDOCARDIUM	(20)	(32)	(30)
HYPERPLASIA, NOS		1 (3%)	
DIGESTIVE SYSTEM			
#LIVER	(20)	(46)	(47)
INFLAMMATION, CHRONIC		1 (2%)	
METAMORPHOSIS FATTY		1 (2%)	
FOCAL CELLULAR CHANGE			1 (2%)
*BILE DUCT	(20)	(50)	(50)
HYPERPLASIA, NOS		1 (2%)	

NUMBER OF ANIMALS WITH TISSUE EXAMINED MICROSCOPICALLY

* NUMBER OF ANIMALS NECROPSIED

**EXCLUDES PARTIALLY AUTOLYZED ANIMALS

TABLE C2 (CONTINUED)

	CONTROL (VEH) 01-F023	LOW DOSE 01-F026	HIGH DOSE 01-F027
#STOMACH ULCER, FOCAL	(20) 2 (10%)	(31)	(31)
URINARY SYSTEM			
#KIDNEY HYDRONEPHROSIS	(20)	(32) 1 (3%)	(30)
INFLAMMATION, SUPPURATIVE	1 (5%)		
INFLAMMATION, CHRONIC	3 (15%)	1 (3%)	5 (17%)
CALCIUM DEPOSIT			1 (3%)
ENDOCRINE SYSTEM			
#ADRENAL ANGIECTASIS	(20)	(30) 1 (3%)	(31) 1 (3%)
#PARATHYROID HYPERPLASIA, NOS	(19) 1 (5%)	(28)	(31)
REPRODUCTIVE SYSTEM			
*VAGINA INFLAMMATION, NOS	(20) 2 (10%)	(50)	(50)
POLYP			1 (2%)
#UTERUS HYDROMETRA	(19) 2 (11%)	(34) 3 (9%)	(35) 4 (11%)
CYST, NOS			1 (3%)
INFLAMMATION, NOS	1 (5%)	3 (9%)	
#CERVIX UTERI EPIDERMAL INCLUSION CYST	(19)	(34)	(35) 1 (3%)
INFLAMMATION, NOS		1 (3%)	
#UTERUS/ENDOMETRIUM HYPERPLASIA, CYSTIC	(19)	(34) 5 (15%)	(35) 2 (6%)
NERVOUS SYSTEM			
NONE			

NUMBER OF ANIMALS WITH TISSUE EXAMINED MICROSCOPICALLY
 * NUMBER OF ANIMALS NECROPSIED

TABLE C2 (CONCLUDED)

	CONTROL (VEH) 01-F023	LOW DOSE 01-F026	HIGH DOSE 01-F027
SPECIAL SENSE ORGANS			
*EYE	(20)	(50)	(50)
SYNECHIA, POSTERIOR			1 (2%)
CATARACT		2 (4%)	2 (4%)
CALCIUM DEPOSIT			1 (2%)
*EYE/RETINA	(20)	(50)	(50)
INFLAMMATION, NOS		1 (2%)	
MUSCULOSKELETAL SYSTEM			
NONE			
BODY CAVITIES			
*ABDOMINAL CAVITY	(20)	(50)	(50)
NECROSIS, FAT		1 (2%)	
*MESENTERY	(20)	(50)	(50)
NECROSIS, FAT			1 (2%)
ALL OTHER SYSTEMS			
NONE			
SPECIAL MORPHOLOGY SUMMARY			
NO LESION REPORTED	5	12	12
NECROPSY PERF/NO HISTO PERFORMED		3	1
# NUMBER OF ANIMALS WITH TISSUE EXAMINED MICROSCOPICALLY			
* NUMBER OF ANIMALS NECROPSIED			

APPENDIX D

SUMMARY OF THE INCIDENCE OF NONNEOPLASTIC
LESIONS IN MICE TREATED WITH METHOXYCHLOR

TABLE DI
SUMMARY OF THE INCIDENCE OF NONNEOPLASTIC LESIONS IN MALE MICE TREATED WITH METHOXYCHLOR

	CONTROL (VEH) 02-M024	LOW DOSE 02-M025	HIGH DOSE 02-M026
ANIMALS INITIALLY IN STUDY	20	50	50
ANIMALS MISSING			1
ANIMALS NECROPSIED	12	44	47
ANIMALS EXAMINED HISTOPATHOLOGICALLY**	12	35	31
INTEGUMENTARY SYSTEM			
*SKIN	(12)	(44)	(47)
INFLAMMATION, NOS		2 (5%)	
*SUBCUT TISSUE	(12)	(44)	(47)
ABSCCESS, NOS			3 (6%)
RESPIRATORY SYSTEM			
#LUNG	(13)	(36)	(33)
PNEUMONIA, CHRONIC MURINE		2 (6%)	
HEMATOPCIIETIC SYSTEM			
*SPLEEN	(12)	(36)	(33)
AMYLOIDOSIS	4 (33%)	9 (25%)	2 (6%)
HEMATOPOIESIS		1 (3%)	1 (3%)
*MESENTERIC L. NODE	(13)	(36)	(31)
CONGESTION, NOS	1 (8%)		
EDEMA, NOS	1 (8%)		
CIRCULATORY SYSTEM			
*HEART	(13)	(36)	(33)
THROMBOSIS, NOS		2 (6%)	
*MYOCARDIUM	(13)	(36)	(33)
INFLAMMATION, NOS		1 (3%)	
DEGENERATION, NOS		1 (3%)	
*AORTA	(12)	(44)	(47)
THROMBUS, ORGANIZED		1 (2%)	

NUMBER OF ANIMALS WITH TISSUE EXAMINED MICROSCOPICALLY

* NUMBER OF ANIMALS NECROPSIED

**EXCLUDES PARTIALLY AUTOLYZED ANIMALS

TABLE D1 (CONTINUED)

	CONTROL (VEH) 02-M024	LOW DOSE 02-M025	HIGH DOSE 02-M026
DIGESTIVE SYSTEM			
#LIVER	(13)	(36)	(32)
INFLAMMATION, NOS		1 (3%)	
METAMORPHOSIS FATTY	1 (8%)		
URINARY SYSTEM			
#KIDNEY	(12)	(36)	(33)
HYDRONEPHROSIS	1 (8%)	1 (3%)	
CYST, NOS			1 (3%)
PYELONEPHRITIS, NOS	1 (8%)	2 (6%)	1 (3%)
INFLAMMATION, CHRONIC	3 (25%)	8 (22%)	1 (3%)
AMYLOIDOSIS	1 (8%)	4 (11%)	
CALCIUM DEPOSIT		2 (6%)	
#URINARY BLADDER	(11)	(35)	(33)
INFLAMMATION, NOS		1 (3%)	
ENDOCRINE SYSTEM			
NONE			
REPRODUCTIVE SYSTEM			
*PREPUTIAL GLAND	(12)	(44)	(47)
INFLAMMATION, NOS	1 (8%)		
*SEMINAL VESICLE	(12)	(44)	(47)
INFLAMMATION, NOS		1 (2%)	
ATROPHY, NOS		1 (2%)	
#TESTIS	(11)	(35)	(32)
CALCIUM DEPOSIT		1 (3%)	
ATROPHY, NOS		3 (9%)	
*EPIDIDYMI	(12)	(44)	(47)
INFLAMMATION, NOS			1 (2%)
GRANULOMA, SPERMATIC			5 (11%)
*SCROTUM	(12)	(44)	(47)
INFLAMMATION, NOS		2 (5%)	
NERVOUS SYSTEM			
NONE			
# NUMBER OF ANIMALS WITH TISSUE EXAMINED MICROSCOPICALLY			
* NUMBER OF ANIMALS NECROPSIED			

TABLE D1 (CONCLUDED)

	CONTROL (VEH) 02-M024	LOW DOSE 02-M025	HIGH DOSE 02-M026
SPECIAL SENSE ORGANS			
NONE			
MUSCULOSKELETAL SYSTEM			
NONE			
BODY CAVITIES			
NONE			
ALL OTHER SYSTEMS			
NONE			
SPECIAL MORPHOLOGY SUMMARY			
NO LESION REPORTED	4	11	13
ANIMAL MISSING/NO NECROPSY			1
NECROPSY PERF/NO HISTO PERFORMED		9	16
AUTO/NECROPSY/HISTO PERF		4	
AUTOLYSIS/NO NECROPSY	8	6	2

TABLE D2
SUMMARY OF THE INCIDENCE OF NONNEOPLASTIC LESIONS IN FEMALE MICE TREATED WITH METHOXYCHLOR

	CONTROL (VEH) 02-F024	LOW DOSE 02-F027	HIGH DOSE 02-F028
ANIMALS INITIALLY IN STUDY	20	50	50
ANIMALS MISSING		1	
ANIMALS NECROPSIED	20	46	50
ANIMALS EXAMINED HISTOPATHOLOGICALLY**	20	16	14
INTEGUMENTARY SYSTEM			
NONE			
RESPIRATORY SYSTEM			
NONE			
HEMATOPOIETIC SYSTEM			
NONE			
CIRCULATORY SYSTEM			
NONE			
DIGESTIVE SYSTEM			
#STOMACH	(20)	(17)	(14)
INFLAMMATION, NOS			1 (7%)
URINARY SYSTEM			
NONE			
ENDOCRINE SYSTEM			
NONE			
REPRODUCTIVE SYSTEM			
#UTERUS	(20)	(17)	(13)
HYPERMETRA	4 (20%)	1 (6%)	
INFLAMMATION, NOS		1 (6%)	
# NUMBER OF ANIMALS WITH TISSUE EXAMINED MICROSCOPICALLY			
* NUMBER OF ANIMALS NECROPSIED			
**EXCLUDES PARTIALLY AUTOLYZED ANIMALS			

TABLE D2 (CONCLUDED)

	CONTROL (VEH) 02-FC24	LOW DOSE 02-FC27	HIGH DOSE 02-FC28
#UTERUS/ENDOMETRIUM HYPERPLASIA, CYSTIC	(20) 3 (15%)	(17) 4 (24%)	(13)
#OVARY CYST, NOS INFLAMMATION, NOS	(20) 5 (25%)	(17) 2 (12%) 1 (6%)	(14)
NERVOUS SYSTEM			
NCNE			
SPECIAL SENSE ORGANS			
*EYE INFLAMMATION, NOS CATARACT	(20) 1 (5%) 1 (5%)	(46)	(50)
MUSCULOSKELETAL SYSTEM			
NCNE			
BODY CAVITIES			
NONE			
ALL OTHER SYSTEMS			
NONE			
SPECIAL MORPHOLOGY SUMMARY			
NO LESION REPORTED	6	6	10
ANIMAL MISSING/NO NECROPSY		1	
NECROPSY PERF/NO HISTO PERFORMED		30	36
AUTOLYSIS/NO NECROPSY		3	
# NUMBER OF ANIMALS WITH TISSUE EXAMINED MICROSCOPICALLY			
* NUMBER OF ANIMALS NECROPSIED			

Review of the Bioassay of Methoxychlor* for Carcinogenicity
by the Data Evaluation/Risk Assessment Subgroup of the
Clearinghouse on Environmental Carcinogens

November 28, 1977

The Clearinghouse on Environmental Carcinogens was established in May, 1976 under the authority of the National Cancer Act of 1971 (P.L. 92-218). The purpose of the Clearinghouse is to advise on the National Cancer Institute's bioassay program to identify and evaluate chemical carcinogens in the environment to which humans may be exposed. The members of the Clearinghouse have been drawn from academia, industry, organized labor, public interest groups, State health officials, and quasi-public health and research organizations. Members have been selected on the basis of their experience in carcinogenesis or related fields and, collectively, provide expertise in organic chemistry, biochemistry, biostatistics, toxicology, pathology, and epidemiology. Representatives of various Governmental agencies participate as ad hoc members. The Data Evaluation/Risk Assessment Subgroup of the Clearinghouse is charged with the responsibility of providing a peer review of NCI bioassay reports on chemicals studied for carcinogenicity. In this context, below is the edited excerpt from the minutes of the Subgroup's meeting at which Methoxychlor was reviewed.

The primary reviewer agreed with the conclusion stated in the report, i.e., that Methoxychlor was not carcinogenic in either the rats or mice under the conditions of test. An experimental shortcoming, mentioned by the reviewer, was that other chemicals were tested at the same time and in the same room as Methoxychlor. He also noted that the treated animals were stressed as a result of the high dosages tested and that the number of matched control animals were inadequate. In conclusion, he opined that Methoxychlor is a safe pesticide with respect to human risk.

A motion was made that the report on the bioassay of Methoxychlor be accepted. The motion was seconded and approved unanimously.

Members present were:

Gerald N. Wogan (Chairman), Massachusetts Institute of
Technology
Lawrence Garfinkel, American Cancer Society
Henry C. Pitot, University of Wisconsin Medical Center
George Roush, Jr., Monsanto Company
Verald K. Rowe, Dow Chemical U.S.A.
Michael B. Shimkin, University of California at San Diego
Louise Strong, University of Texas Health Sciences Center
John H. Weisburger, American Health Foundation

-
- * Subsequent to this review, changes may have been made in the bioassay report either as a result of the review or other reasons. Thus, certain comments and criticisms reflected in the review may no longer be appropriate.



4 0128 2362

NIH LIBRARY



3 1496 00185 2634